

ON $\mathcal{I}^{\mathcal{K}}$ -CONVERGENCE IN TOPOLOGICAL SPACES VIA SEMI-OPEN SETS

Ankur Sharmah and Debajit Hazarika

Abstract. A sequence $\{x_n\}$ is $s\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to ξ , if there exists a ‘big enough’ subsequence $\{x_{n_k}\}$ which \mathcal{K} -converges to ξ via semi-open sets. In this paper, we introduce the concept of $s\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence which generalizes $\mathcal{S}\text{-}\mathcal{I}$ -convergence and discuss some properties, as well as its relation with compact sets. For two given ideals \mathcal{I} and \mathcal{K} , we justify the existence of an ideal such that $\mathcal{I}^{\mathcal{K}}$ -convergence and convergence with the third ideal coincides for semi-open sets. Moreover, the notion of $s\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point of a sequence is defined and studied here. We characterize the collection of $s\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster points of a sequence as semi-closed subsets of the space.

1. Introduction

After Kuratowski introduced ideals in 1933, the term became known as a collection of sets considered to be “small” or “negligible”. In an ordinary space, three basic topological notions, namely convergence, closure and neighborhood, play a crucial role in determining other topological properties. In the recent past, ideal theory has been used together with convergence theory to develop some promising generalizations of existing concepts in Point-Set Topology.

In particular, two notions for the convergence of a sequence were introduced in 2000 by Kostyrko et al. [7], called \mathcal{I} and \mathcal{I}^* -convergence for the real numbers and later, in 2005, by Lahiri and Das [8] for a topological space. Undoubtedly, it was actively practiced in the following period, and some work from it can be found in [4]. Later, in 2011, Macaz and Sleziak [10] introduced the notion of $\mathcal{I}^{\mathcal{K}}$ -convergence of a function in a topological space. Although it appears in the context of \mathcal{I}^* -convergence, $\mathcal{I}^{\mathcal{K}}$ -convergence further extends the notion of ideal convergence. In particular, if the ideals \mathcal{I} and \mathcal{K} coincide, then the terms $\mathcal{I}^{\mathcal{K}}$ and \mathcal{I} -convergence also apply. In the last decade, $\mathcal{I}^{\mathcal{K}}$ -convergence has been studied in detail in several articles, namely, [3, 10].

2020 Mathematics Subject Classification: 40A35, 40A05, 54A05, 54A20

Keywords and phrases: $\mathcal{I}^{\mathcal{K}}$ -convergence; $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence; $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster points; semi-open sets; semi-compactness; semi-dense set.

On the other hand, Levine introduced semi-open sets [9] in a topological space in 1963, and afterwards it was used to generalize several concepts in Point-Set Topology. Recently, Guevara et al. [5] used the concept of semi-open set to define and study the notion of \mathcal{S} - \mathcal{I} -convergence in topological spaces. In this paper, we define $\mathcal{I}^{\mathcal{K}}$ -convergence using semi-open sets in a topological space and denote it by \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -convergence.

An ideal on a set S is a collection of subsets of the given set that is closed under subset inclusion and finite union. Fin is a basic ideal that includes all finite subsets of S . For a given ideal $\mathcal{I} \subset P(\mathbb{N})$, two additional subsets of $P(\mathbb{N})$ namely, \mathcal{I}^* or $F(\mathcal{I})$ and \mathcal{I}^+ of $P(\mathbb{N})$, are defined, namely a: $\mathcal{I}^* := \{A \subset \mathbb{N} : A^c \in \mathcal{I}\}$ and $\mathcal{I}^+ :=$ collection of all subsets that do not belong to \mathcal{I} . We say that two ideals \mathcal{I} and \mathcal{K} on S fulfill the ideality condition if $\mathcal{I} \cup \mathcal{K}$ is a proper ideal [14], alternatively, $S \neq I \cup K$, for all $I \in \mathcal{I}, K \in \mathcal{K}$.

In this paper, we deal with the proper ideals (not containing \mathbb{N}) on the set of natural numbers \mathbb{N} , which are admissible ideals (containing all finite subsets of \mathbb{N}), to study different aspects of \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -convergence in a topological space. Thus, in the following part of this paper, all considered ideals are proper and admissible. The main results in this paper are divided into two sections.

Section 2 introduces the \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -convergence for two ideals \mathcal{I}, \mathcal{K} satisfying the ideality condition and some basic properties are investigated. Section 4 deals with the definition and basic properties of \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -cluster points of a sequence. The two sections 3 and 5 contribute to answering an existence problem (Theorem 3.14, Theorem 5.5), which can be formulated as follows: Whether there exists an ideal \mathcal{J} for two given ideals \mathcal{I}, \mathcal{K} such that \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -convergence and \mathcal{S} - \mathcal{J} -convergence coincide under certain assumptions. Section 5 also contains the characterization of the collection $sC_x(\mathcal{I}^{\mathcal{K}})$ of \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -cluster points of a sequence as semi-closed subsets (Theorem 5.8) and further, we evaluate a condition for coincidence for the collection of $sC_x(\mathcal{I}^{\mathcal{K}})$ and $sL(\mathcal{I}^{\mathcal{K}})$ (Theorem 5.3).

2. Preliminaries

For a given function $f : S \rightarrow X$, which is in fact a generalization of a sequence, Macaz and Sleziaak [10] defined the $\mathcal{I}^{\mathcal{K}}$ -convergence for two ideals \mathcal{I} and \mathcal{K} on S .

DEFINITION 2.1 ([10]). A function $f : S \rightarrow X$ is said to be \mathcal{K} -convergent to $x \in X$, if for any nonempty open set U containing x , we have $\{s \in S : f(s) \notin U\} \in \mathcal{K}$.

DEFINITION 2.2 ([10]). A function $f : S \rightarrow X$ is said to be $\mathcal{I}^{\mathcal{K}}$ -convergent to $x \in X$, if there exists a set $M \in F(\mathcal{I})$ such that the function $g : S \rightarrow X$ given by $g(s) = f(s)$, if $s \in M$ and $g(s) = x$, if $s \notin M$, is \mathcal{K} -convergent to x . If f is $\mathcal{I}^{\mathcal{K}}$ -convergent to x , then we write $\mathcal{I}^{\mathcal{K}}\text{-lim } f = x$.

In particular, following is the definition of $\mathcal{I}^{\mathcal{K}}$ -convergence of a sequence in a space.

DEFINITION 2.3. A sequence $\{x_n\}$ is said to be $\mathcal{I}^{\mathcal{K}}$ -convergent to an element $\xi \in X$ if there exists a set $M = \{n_1, n_2, \dots, n_k, \dots\} \in \mathcal{I}^*$ such that the subsequence $\{x_{n_k}\}$ is \mathcal{K} -convergent to ξ .

PROPOSITION 2.4 ([14, Proposition 2.1]). *Let X be a topological space and $f : S \rightarrow X$ be a function. Let \mathcal{I}, \mathcal{K} be two ideals on S such that $\mathcal{I} \cup \mathcal{K}$ is an ideal. Then*

(i) $\mathcal{I}^{\mathcal{K}^*}$ - $\lim f = x$ if and only if $(\mathcal{I} \cup \mathcal{K})^*$ - $\lim f = x$.

(ii) $\mathcal{I}^{\mathcal{K}}$ - $\lim f = x$ implies $\mathcal{I} \cup \mathcal{K}$ - $\lim f = x$.

Some of the definitions and concepts of generalized open sets in [2, 5, 9, 11, 12] that are used in the content of the accompanying sections are listed below.

DEFINITION 2.5. Let X be a topological space. Then

(i) $O \subset X$ is said to be semi-open [9] if there exists an open set U such that $U \subset O \subset \bar{U}$. The collection of all semi-open subsets of X is denoted by $SO(X)$.

(ii) The complement set of a semi-open set is termed as a semi-closed set.

(iii) The semi-closure [9] of a subset F of X , denoted by $sCl(F)$, is defined as the intersection of all semi-closed set containing F . Otherwise, a point $x \in sCl(A)$ if and only if for every semi-open set U containing x , $U \cap A \neq \emptyset$.

(iv) An element $x \in F \subset X$ is said to be semi-limit point [2] of F , if for every semi-open set O containing x , $O \cap F \neq \phi$.

(v) A topological space X is said to be semi-Hausdorff [11] if for every distinct pair of elements $x, y \in X$, there exists a disjoint pair of semi-open sets U and V containing x and y respectively.

(vi) A function $f : X \rightarrow Y$ is said to be irresolute [5] if $f^{-1}(O) \in SO(X)$ for each $O \in SO(Y)$. A function $f : X \rightarrow Y$ is irresolute [5] if and only if for each $x \in X$ and each $V \in SO(Y)$ containing $f(x)$, there exists $U \in SO(X)$ such that $x \in U$ and $f(U) \subset V$.

(vii) A function $f : X \rightarrow Y$ is said to be semi-continuous [9] if $f^{-1}(O) \in SO(X)$ for each open set $O \in Y$. A function $f : X \rightarrow Y$ is semi-continuous [9] if and only if for each $x \in X$ and each open V in Y containing $f(x)$, there exists $U \in SO(X)$ such that $x \in U$ and $f(U) \subset V$.

THEOREM 2.6 ([5, Theorem 3.5]). *Let X be a space and \mathcal{I} be an ideal. If every sequence $\{x_n\}$ in X has an \mathcal{S} - \mathcal{I} -cluster point, then every infinite subset of X has a semi- ω -accumulation point. The converse is true if \mathcal{I} does not contain any infinite sets.*

Meanwhile, we refer to [6, 8] for the basic general topological and ideal theoretic terminologies, definitions and results mentioned in the content. In the following, unless otherwise stated, we denote X as topological space and \mathcal{I} and \mathcal{K} as ideals on \mathbb{N} .

3. Some properties of $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence

As a generalization of the work on $\mathcal{S}\text{-}\mathcal{I}$ -convergence, we consider $\mathcal{I}^{\mathcal{K}}$ -convergence, which is one of the most generalized among all ideal convergences, to study its properties using semi-open sets. We denote it as $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence in a space X . A sequence $x = \{x_n\}$ is called $\mathcal{S}\text{-}\mathcal{I}$ -convergent [5] to an element $\xi \in X$ if for every non-empty semi-open set U containing ξ , the set $\{n \in \mathbb{N} : x_n \notin U\}$ belongs to \mathcal{I} . Here we define the notion of $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence of a sequence in a topological space.

DEFINITION 3.1. A sequence $\{x_n\}$ is said to be $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{M}}$ -convergent to an element $\xi \in X$ if there exists a set $M = \{n_1, n_2, \dots, n_k, \dots\} \in \mathcal{I}^*$ such that the subsequence $\{x_{n_k}\}$ is $\mathcal{S}\text{-}\mathcal{M}$ -convergent to ξ , where \mathcal{M} is an ideal convergence mode.

If \mathcal{K} is an ideal and $\mathcal{M} = \mathcal{K}^*$, then we say that $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ -convergent to an element $\xi \in X$. Also, if $\mathcal{M} = \mathcal{K}$, then $\{x_n\}$ is said to be $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to an element $\xi \in X$. If the ideal \mathcal{K} does not contain an infinite set, then Definition 3.1 exhibits the $\mathcal{S}\text{-}\mathcal{I}^*$ -convergence. Also, if \mathcal{K} is a P-ideal [3] (condition AP [8]), then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ and $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence both coincide. Basically, \mathcal{K} -convergence implies $\mathcal{I}^{\mathcal{K}}$ -convergence (Definition 2.3), analogously we have the following lemma.

LEMMA 3.2. $\mathcal{S}\text{-}\mathcal{K}$ convergence implies $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence.

Proof. Let $\{x_n\}$ be an $\mathcal{S}\text{-}\mathcal{K}$ -convergent sequence to an element ξ in a space X . Then, for any semi-open set U containing ξ , we have $\{n \in \mathbb{N} : x_n \notin U\} \in \mathcal{K}$. Then, for any $M \in \mathcal{I}^*$, the set $\{n \in M : x_n \notin U\} \subset \{n \in \mathbb{N} : x_n \notin U\} \in \mathcal{K}$. Thus, $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent in X . \square

PROPOSITION 3.3. Let $\{x_n\}$ be a sequence in X . For \mathcal{I}, \mathcal{K} be two ideals on \mathbb{N} such that $\mathcal{I} \cup \mathcal{K}$ is an ideal. Then

- (i) $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ -converges to x if and only if $\{x_n\}$ is $\mathcal{S}\text{-}(\mathcal{I} \cup \mathcal{K})^*$ -converges to x .
- (ii) Also, $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -converges to x implies $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I} \cup \mathcal{K}$ -converges to x .

Proof. Consider two ideals \mathcal{I} and \mathcal{K} such that $\mathcal{I} \cup \mathcal{K}$ is an ideal.

(i) Let $\{x_n\}$ be a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ -convergent to x . So, there exists $M = \{n_1, n_2, n_3, \dots\}$ such that $\{x_{n_k}\}$ is $\mathcal{S}\text{-}\mathcal{K}^*$ -convergent to x . Then, for any semi open set U containing x , there exists $N \in \mathcal{K}^*$ such that $\{n_k \in N : x_{n_k} \notin U\} \in Fin$, i.e., $\{n \in M \cap N : x_n \notin U\} \in Fin$. But $M \cap N \in (\mathcal{I} \cap \mathcal{K})^*$. Hence, $\{x_n\}$ is $\mathcal{S}\text{-}(\mathcal{I} \cup \mathcal{K})^*$ -convergent to x .

Conversely, let $\{x_n\}$ be $\mathcal{S}\text{-}(\mathcal{I} \cup \mathcal{K})^*$ -convergent to x . So, for a semi-open set U containing x , there exists $M \cap N \in (\mathcal{I} \cup \mathcal{K})^*$, where $M \in \mathcal{I}^*, N \in \mathcal{K}^*$, such that $\{n \in M \cap N : x_n \notin U\} \in Fin$. Consider a subsequence $\{x_{n_k}\}_{n_k \in M}$ of $\{x_n\}_{n \in \mathbb{N}}$ such that $\{n_k \in N : x_{n_k} \notin U\} \in Fin$. Thus, $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ -convergent to x .

(ii) Let $\{x_n\}$ be $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to x . So, there exists $M \in \mathcal{I}^*$ such that $\{n \in M : x_n \notin U_x\} \in \mathcal{K}$, where U is an semi open set containing x . Then, $\{n \in \mathbb{N} : x_n \notin U\} \subset \{n \in M : x_n \notin U\} \cup \{n : n \notin M\}$. Therefore, $\{n \in \mathbb{N} : x_n \notin U\} \in \mathcal{I} \cup \mathcal{K}$. Thus, $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I} \cup \mathcal{K}$ -convergent to x . \square

Following results are immediate consequences of Proposition 3.3.

COROLLARY 3.4. Let \mathcal{I} and \mathcal{K} be two ideals on \mathbb{N} provided $\mathcal{I} \cup \mathcal{K}$ is an ideal. Then

(i) $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ -convergence implies both $\mathcal{S}\text{-}\mathcal{I}$ -convergence as well as $\mathcal{S}\text{-}\mathcal{K}$ -convergence.

(ii) $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence implies $\mathcal{S}\text{-}\mathcal{I}$ -convergence provided $\mathcal{K} \subseteq \mathcal{I}$.

(iii) $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence implies $\mathcal{S}\text{-}\mathcal{K}$ -convergence provided $\mathcal{I} \subseteq \mathcal{K}$.

If $\mathcal{K} = \text{Fin}$, then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence implies $\mathcal{S}\text{-}\mathcal{I}$ -convergence. A simple observation is that if the ideals \mathcal{I} and \mathcal{K} both coincide, then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence appear as the $\mathcal{S}\text{-}\mathcal{I}$ -convergence. Eventually, if $\mathcal{I}, \mathcal{K} = \text{Fin}$, then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence coincide with \mathcal{S} -convergence and hence, further implies usual convergence. Following is an example to show that even if $\mathcal{I} = \mathcal{K} = \text{Fin}$, usual convergence does not coincide with the $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence.

EXAMPLE 3.5. Let \mathcal{I} and \mathcal{K} be two ideals in \mathbb{N} such that $\mathcal{I} \cup \mathcal{K}$ is an ideal. Let \mathbb{R} be the set of real numbers with usual topology and let $\{x_n\}$ be defined as $x_n = (\frac{1}{n})$. Then $x_n \rightarrow \text{is}0$. Consider the semi-open set $U = (-1, 0]$ containing 0. But $\{n \in \mathbb{N} : x_n \notin U\} = \mathbb{N} \notin \mathcal{I} \cup \mathcal{K}$ for any ideals \mathcal{I} and \mathcal{K} . So $\{x_n\}$ is not $\mathcal{S}\text{-}\mathcal{I} \cup \mathcal{K}$ -convergent to 0. Therefore, $\{x_n\}$ is not $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to 0 by Proposition 3.3.

The proof of the next lemma follows immediately from the definition of $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence.

LEMMA 3.6. $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence implies $\mathcal{I}^{\mathcal{K}}$ -convergence, for any ideals \mathcal{I} and \mathcal{K} .

The following example shows that the converse of Lemma 3.6 is not necessarily true.

EXAMPLE 3.7. Let \mathcal{I} and \mathcal{K} be two ideals in \mathbb{N} such that $\mathcal{I} \cup \mathcal{K}$ is an ideal. Let $[-1, 1]$ be the interval in \mathbb{R} with usual subspace topology and $\{x_n\}$ a sequence defined as $x_n = (\frac{1}{n}) \sin(\frac{1}{n})$. Thus, for any open set U containing 0, we have $\{n \in \mathbb{N} : x_n \notin U\}$ is finite, that implies $x_n \rightarrow_{\mathcal{K}} 0$. Therefore, $\{x_n\}$ is $\mathcal{I}^{\mathcal{K}}$ -convergent to 0. Let us now consider the semi-open set $V = (-1, 0]$ in $[-1, 1]$. Then $\{n \in \mathbb{N} : x_n \notin V\} = \mathbb{N} \notin \mathcal{I} \cup \mathcal{K}$. Hence, $x_n \not\rightarrow_{\mathcal{I} \cup \mathcal{K}} 0$. It therefore follows from Proposition 2.4 (ii) that $\{x_n\}$ is not $\mathcal{I}^{\mathcal{K}}$ -convergent to 0.

THEOREM 3.8. In a semi-Hausdorff space X , each $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent sequence has a unique $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit in X , provided $\mathcal{I} \cup \mathcal{K}$ is an ideal.

Proof. Consider a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent sequence $\{x_n\}$ in a semi-Hausdorff space X . Suppose that $\{x_n\}$ has two distinct $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limits, say a and b . Being X a semi-Hausdorff space, there exists $U, V \in \mathcal{S}O(X)$ with $U \cap V = \emptyset$ such that $a \in U, b \in V$. As $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to a and b , there exist $M_1, M_2 \in \mathcal{I}^*$ such that $\{n \in M_1 : x_n \notin U\} \in \mathcal{K}$ and $\{n \in M_2 : x_n \notin V\} \in \mathcal{K}$. So, for $M = M_1 \cap M_2 \in \mathcal{I}^*$ ($\neq \emptyset$), the sets $\{n \in M : x_n \notin U\}$ and $\{n \in M : x_n \notin V\}$ belong to \mathcal{K} . Now, we have

$$\{n \in M : x_n \notin U \cap V\} = \{n \in M : x_n \notin U\} \cup \{n \in M : x_n \notin V\} \in \mathcal{K}.$$

Again,

$$\{n \in \mathbb{N} : x_n \notin U \cap V\} = \{n \in M : x_n \notin U \cap V\} \cup \{n \in M^c : x_n \notin U \cap V\}$$

$$\subseteq \{n \in M : x_n \notin U \cap V\} \cup M^{\mathcal{G}} \in \mathcal{I} \cup \mathcal{K}.$$

But, $\mathcal{I} \cup \mathcal{K}$ is an ideal, which implies $\{n \in \mathbb{N} : x_n \notin U \cap V\} \neq \mathbb{N}$. Therefore, we conclude that $\{n \in \mathbb{N} : x_n \in U \cap V\} \neq \emptyset$, which is a contradiction. \square

COROLLARY 3.9. *In a Hausdorff space X , each $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent sequence has a unique $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit in X , provided $\mathcal{I} \cup \mathcal{K}$ is an ideal.*

REMARK 3.10. Since sequences in a general space are inadequate, the unique $\mathcal{I}^{\mathcal{K}}$ -limit of every X -valued sequence does not necessarily imply that X is Hausdorff. But the space X will be at least T_1 . If not, then there exists one distinct pair $x, y \in X$ such that for U (open set) containing x , it also contains y , or for U (open set) containing y , it also contains x . Without loss of generality, we assume that for U (open set) containing x implies that $y \in U$. Consider the sequence $\{x_n\}$ such that $x_n = y, \forall n \in \mathbb{N}$. Undoubtedly, $\{x_n\}$ is $\mathcal{I}^{\mathcal{K}}$ -convergent to y . However, since x shares all its open sets with y , every sequence $\mathcal{I}^{\mathcal{K}}$ that converges to y also converges to x . Therefore, $\{x_n\}$ is $\mathcal{I}^{\mathcal{K}}$ -convergent to x . This contradicts our assumption. Thus X is a T_1 -space. If X is a first countable space, then X is also Hausdorff.

Immediately, we observe that the above remark is true for $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence in a semi-Hausdorff space. So, using Theorem 3.8, we have the following result.

THEOREM 3.11. *If each sequence in a space X has a unique $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit, then X has the semi- T_1 [11] property. Moreover, X is a semi-Hausdorff space provided X is first countable.*

In a space, uniqueness of $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit of sequences is a stronger property than that of usual limit. So, following question is relevant in this context: **Does the uniqueness of $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit of every sequence in X imply that the space X is Hausdorff?**

Proposition 3.3 hints at an interesting existence scenario which can be stated as whether there exists an ideal \mathcal{J} such that $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence coincide with $\mathcal{S}\text{-}\mathcal{J}$ -convergence. Similar results for $\mathcal{I}^{\mathcal{K}}$ -convergence is affirmatively answered for Hausdorff spaces in article [14]. Following results contribute to the problem of interlinking between $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergence and usual ideal convergence.

REMARK 3.12. If \mathcal{I} and \mathcal{K} be two ideals on \mathbb{N} satisfying the ideality condition. Consider the set $\{K \cup J : K \in \mathcal{K}\}$, for any $J \in \mathcal{I}$. If \mathcal{J} is the collections of sets under the operations finite union and subset inclusion among the members of $\{K \cup J : K \in \mathcal{K}\}$. Then, \mathcal{J} is an ideal on \mathbb{N} . Also, $\mathcal{J} \subseteq \mathcal{I} \cup \mathcal{K}$.

LEMMA 3.13. *Let \mathcal{I}, \mathcal{K} be two ideals on \mathbb{N} satisfying ideality condition. Let $\{x_n\}$ be a X -valued sequence. If $\mathcal{J} =$ ideal generated by the set $\{K \cup J : K \in \mathcal{K}\}$, for any $J \in \mathcal{I}$. Then $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{J}$ -convergent to x implies $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to x .*

Proof. Let $x = \{x_n\}$ be $\mathcal{S}\text{-}\mathcal{J}$ -convergent in X , where $\mathcal{J} =$ ideal generated by the set $\{K \cup J : K \in \mathcal{K}\}$, for any $J \in \mathcal{I}$. Assuming $M = J^{\mathcal{G}}$ and $V \in \mathcal{S}\mathcal{O}(X)$, we can observe that $\{n_k \in M : x_{n_k} \notin V\} \subseteq \{n \in \mathbb{N} : x_n \notin V\} \setminus \{n_k \in \mathbb{N} : n_k \notin M\}$. Again,

$\{n_k \in M : x_{n_k} \notin V\} \setminus J \subseteq (K \cup J) \setminus J \in \mathcal{K}$. Subsequently, x_{n_k} is $\mathcal{S}\text{-}\mathcal{K}$ -convergent to x . Hence, $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to x . \square

THEOREM 3.14. *Let X be a Hausdorff Space. Let $\{x_n\}$ be $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to x . Then, there exists an ideal \mathcal{J} such that $x \in X$ is an $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit of the sequence $\{x_n\}$ if and only if x is also a \mathcal{J} -limit of $\{x_n\}$ provided $\mathcal{I} \cup \mathcal{K}$ is an ideal.*

Proof. Suppose that $\{x_n\}$ is $\mathcal{I}^{\mathcal{K}}$ -convergent to x . So, there exists a set $M \in \mathcal{I}^*$ such that $\{n_k \in M : x_{n_k} \notin U_x\} \in \mathcal{K}$ where $U_x \in \mathcal{S}\mathcal{O}(X)$. Now, let $J = M^c$. Since, $(\mathcal{I} \cup \mathcal{K})$ is an ideal, the set $\{K \cup J : K \in \mathcal{K}\}$ generates an ideal, say \mathcal{J} . Then

$$\{n \in \mathbb{N} : x_n \notin U_x\} \subseteq \{n_k \in M : x_{n_k} \notin U_x\} \cup M^c \in \mathcal{J}.$$

Hence, $\{x_n\}$ is \mathcal{J} -convergent to x .

Conversely, suppose that $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{J}$ -convergent to x , where \mathcal{J} = ideal generated by $(\mathcal{K} \cup J)$, for any $J \in \mathcal{I}$. Then, by Lemma 3.13, $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to x . \square

THEOREM 3.15. *Let \mathcal{I} and \mathcal{K} be two ideals and $F \subset X$. If there exists a sequence $\{x_n\}$ in F (with distinct elements) which is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $\xi \in X$, then ξ is a semi-limit point of F , in essence, $\xi \in sCl(F)$, the semi-closure of F .*

Proof. Let U be any semi-open subset of X containing the point ξ . Since $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $\xi \in X$, so, there exists a set $M \in \mathcal{I}^*$ such that $\{n \in M : x_n \notin U\} \in \mathcal{K}$. In other words, $\{n \in M : x_n \in U\} \notin \mathcal{K}$ since \mathcal{K} is an ideal. Then choose $n_0 \in \{n \in M : x_n \in U\}$ such that $x_{n_0} \neq \xi$, then $x_{n_0} \in F \cap (U - \{\xi\})$ and hence, $F \cap (U - \{\xi\}) \neq \emptyset$. This shows that ξ is a semi-limit point of F . \square

COROLLARY 3.16. *Let \mathcal{I} and \mathcal{K} be two ideals and consider $F \subset X$. If there exists a sequence $\{x_n\}$ in F (with distinct elements) which is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $\xi \in X$, then $\xi \in Cl(F)$.*

THEOREM 3.17. *If $F \subset X$ is a semi-closed set, then for any sequence in F which is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to a , we have $a \in F$.*

Proof. Suppose $F \subset X$ is a semi-closed set and $\{x_n\}$ is any sequence in F that is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to the element a , but $a \notin F$. Since F is semi-closed, we have $sCl(F) = F$ and therefore $a \notin sCl(F)$. Then there exists a semi-open set U containing a such that $F \cap U \neq \emptyset$. Since $\{x_n\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to a , there exists $M \in \mathcal{I}^*$ such that $\{n \in M : x_n \notin U\} \in \mathcal{K}$. Furthermore, $\{n \in M : x_n \in U\} \notin \mathcal{K}$, which implies that $\{n \in M : x_n \in U\} \neq \emptyset$. According to our hypothesis, $x_n \in F$, that implies $F \cap U \neq \emptyset$. This is a contradiction. \square

THEOREM 3.18. *Let $f : X \rightarrow Y$ be a semi-continuous function. If $\{x_n\}$ is a sequence in X which is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $\xi \in X$, then $\{f(x_n)\}$ is an $\mathcal{I}^{\mathcal{K}}$ -convergent sequence to $f(\xi)$.*

Proof. Consider a sequence $\{x_n\}$ in X which is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $\xi \in X$. We claim that $\{f(x_n)\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $f(\xi)$. Suppose not, that is $\{f(x_n)\}$ is not $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $f(\xi)$. Then there exists an open set $V \subseteq Y$, containing $f(\xi)$ and for each

$M \in \mathcal{I}^*$, we have $\{n \in M : f(x_n) \notin V\} \notin \mathcal{K}$. Now by Definition 2.5 (vii), there exists $U \in SO(X)$ such that $\xi \in U$ and $f(U) \subset V$. Now, $\{n \in M : f(x_n) \notin V\} \subset \{n \in M : x_n \notin U\}$. Then $\{n \in M : x_n \notin U\} \notin \mathcal{K}$, that implies $\{x_n\}$ is not $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to ξ . That is a contradiction to our assumption. Hence, $\{f(x_n)\}$ is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $f(\xi)$. \square

THEOREM 3.19. *Let $f : X \rightarrow Y$ be an irresolute function. If $\{x_n\}$ is a sequence in X which is $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -convergent to $\xi \in X$, then $\{f(x_n)\}$ $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -converges to $f(\xi)$.*

Proof. The proof is similar to that of Theorem 3.18 with the use of the characterization of an irresolute function that is shown in Definition 2.5 (vi). \square

4. $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster points and several properties

In this section we introduce the terminology of $\mathcal{I}^{\mathcal{K}}$ cluster points of a sequence for semi-open sets in a topological space. An element p in a space X is said to be an \mathcal{I}^* -cluster point of a sequence $\{x_n\}$ if there exists $M = \{m_1, m_2, \dots, m_k, \dots\} \in \mathcal{I}^*$ such that, that the subsequence $\{x_{m_k}\}$ has a cluster point p , more precisely, for any open set U containing p , the set $\{n \in \mathbb{N} : x_{m_k} \in U\}$ is an infinite subset of \mathbb{N} .

DEFINITION 4.1. Let X be a space and $\{x_n\}$ be a sequence in X . A point $p \in X$ is called a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{M}}$ -cluster point of $\{x_n\}$ in X if there exists $M \in \mathcal{I}^*$ such that for any $U \in SO(X)$ containing p , we have $\{n \in M : x_n \in U\} \notin \mathcal{M}$, where \mathcal{M} is an ideal convergence mode.

If $\mathcal{M} = \mathcal{K}$ and $\mathcal{M} = \mathcal{K}^*$, then Definition 4.1 refers to the definitions of $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point and $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}$ -cluster point of a sequence correspondingly. It is doubtless that each $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit of a sequence is a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point, but the converse is not always true. But if the ideal \mathcal{K} is a maximal ideal (for any $A \subset \mathbb{N}$ it implies that $A \in \mathcal{K}$ or $A^c \in \mathcal{K}$), then the converse is also true i.e. $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -limit and $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point of a sequence coincide.

REMARK 4.2. Let X be a space and $\{x_n\}$ be a sequence in X . Then

1. If $\mathcal{I} = \mathcal{K}$, then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster points of $\{x_n\}$ coincide with that of $\mathcal{S}\text{-}\mathcal{I}$ -cluster points.
2. If $\mathcal{K} = Fin$, then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster points of $\{x_n\}$ are also $\mathcal{S}\text{-}\mathcal{I}$ -cluster points.

We denote the collection of all $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster points of a sequence $x = \{x_n\}$ in a space by $sC_x(\mathcal{I}^{\mathcal{K}})$. Consequently, from Definition 4.1, it straightaway follows that $sC_x(\mathcal{I}^{\mathcal{K}}) \subseteq sC_x(\mathcal{K})$.

LEMMA 4.3. $sC_x(\mathcal{I} \cup \mathcal{K}) \subseteq sC_x(\mathcal{I}^{\mathcal{K}})$, for two ideals \mathcal{I} and \mathcal{K} such that $\mathcal{I} \cup \mathcal{K}$ is an ideal.

Proof. Let y be not a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point of $x = \{x_n\}$. Then for all $M \in \mathcal{I}^*$ there exists $U \in SO(X)$ containing y such that $\{n \in M : x_n \in U\} \in \mathcal{K}$. Hence, $\{n \in \mathbb{N} : x_n \in U\} \subseteq \{n \in M : x_n \in U\} \cup M^c \in \mathcal{I} \cup \mathcal{K}$. Hence, y is not a $\mathcal{S}\text{-}(\mathcal{I} \cup \mathcal{K})$ -cluster point of x . \square

THEOREM 4.4. *Let $\mathcal{I}, \mathcal{K}, \mathcal{I}_1, \mathcal{I}_2, \mathcal{K}_1$ and \mathcal{K}_2 be ideals on \mathbb{N} . Suppose that $x = \{x_n\}$, $y = \{y_n\}$ are two sequences in a space X . Then*

(i) *If $\mathcal{K}_1 \subset \mathcal{K}_2$, then $sC_x(\mathcal{I}^{\mathcal{K}_2}) \subseteq sC_x(\mathcal{I}^{\mathcal{K}_1})$,*

(ii) *If $\mathcal{I}_1 \subset \mathcal{I}_2$, then $sC_x(\mathcal{I}_1^{\mathcal{K}}) \subseteq sC_x(\mathcal{I}_2^{\mathcal{K}})$*

(iii) *and if $\{n \in \mathbb{N} : x_n \neq y_n\} \in \mathcal{K}$, then $sC_x(\mathcal{I}^{\mathcal{K}}) = sC_y(\mathcal{I}^{\mathcal{K}})$.*

Proof. The proof of (i) and (ii) follows directly from the Definition 4.1.

(iii) Consider, $N = \{n \in \mathbb{N} : x_n \neq y_n\} \in \mathcal{K}$. Suppose that $a \in sC_x(\mathcal{I}^{\mathcal{K}})$, then there exists $M \in \mathcal{I}^*$ such that for each $U_a \in SO(X)$, we have $\{n \in M : x_n \in U_a\} \notin \mathcal{K}$. We negate the possibility of $M \cap N = \emptyset$, as in that case the result follows immediately. Again, $M \cap N \in \mathcal{K}$ and

$$\{n \in M : y_n \in U_a\} = \{n \in M \cap N : y_n \in U_a\} \cup \{n \in M \setminus N : x_n \in U_a\}.$$

However, $\{n \in M \cap N : y_n \in U_a\} \in \mathcal{K}$. Thus, $\{n \in M : y_n \in U_a\} \notin \mathcal{K}$. Thus, $a \in sC_y(\mathcal{I}^{\mathcal{K}})$. Since the sequences x, y are taken arbitrarily, hence $sC_x(\mathcal{I}^{\mathcal{K}}) = sC_y(\mathcal{I}^{\mathcal{K}})$. \square

Recall that an element $p \in X$ is a semi- ω -accumulation point [5] of $A \subset X$ if for every semi-open set U containing p , $U \cap A$ is an infinite set.

THEOREM 4.5. *Let X be a space and \mathcal{I}, \mathcal{K} be two ideals. If every sequence $\{x_n\}$ in X has a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point, then every infinite subset of X has a semi- ω -accumulation point.*

Proof. Let F (infinite) $\subset X$, then there exists a sequence $\{x_n\}$ of distinct points in F . Suppose that every sequence in X has a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point. Let a be a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point of $\{x_n\}$. Then, for $U \in SO(X)$ containing a , we have $\{n \in M : x_n \in U\} \notin \mathcal{K}$. Since as per assumption, $Fin \subset \mathcal{K}$ then the set $\{n \in M : x_n \in U\}$ is infinite. Since $x_n \in F$, $U \cap F$ is an infinite set. Hence a is semi- ω -accumulation point of F . \square

COROLLARY 4.6. *Let X be a space. If every sequence $\{x_n\}$ has a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point, then every infinite subset of X has a ω -accumulation point.*

Recall that a space X is semi-compact [13] if every semi-open cover of X possesses a finite subcover. Similarly, X is said to be semi-Lindelöf if every semi-open cover of X possesses a countable subcover.

THEOREM 4.7. *If X is a semi-Lindelöf space and each sequence in X has an $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point, then X is a semi-compact space.*

Proof. Suppose that X is a semi-Lindelöf space and that each X -valued sequence has a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point. Let $\mathcal{S} = \{S_\lambda : \lambda \in \Lambda\}$ be a semi-open cover of X . So, there exists a countable subcover $\mathcal{S}' = \{S_1, S_2, \dots, S_m, \dots\}$. If possible, consider the sequence $U = \{U_m\}$ such that $U_1 = S_1$ and for each $m > 1$, let $U_m = S_m$, where S_m is the first member of the sequence of U 's such that $S_m \not\subseteq U_1 \cup U_2 \cup \dots \cup U_{m-1}$. Assuming the axiom of choice, consider a sequence $x = \{x_m\}$ such that $x_1 \in U_1$ and for each $m > 1$, let $x_m \in U_m - (U_1 \cup U_2 \cup \dots \cup U_{m-1})$. Now by our hypothesis, $\{x_m\}$ has a $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster

point, say l . Then, there exists j such that $l \in U_j$. Consequently, there exists a set $M \in \mathcal{I}^*$ such that $\{n \in M : x_n \in U_j\} \notin \mathcal{K}$. So, the set $\{n \in M : x_n \in U_j\}$ must be a infinite subset of \mathbb{N} . Thus, there exists $k > j$ such that $k \in \{n \in M : x_n \in U_j\}$; that is $x_k \in U_j$, which is a contradiction. Subsequently, there must exist $m_0 \in \mathbb{N}$ such that the process of induction for $\{U_m\}$ is impossible to continue after $m = m_0$. Therefore, $\{U_1, U_2, \dots, U_{m_0}\}$ is a finite subcover of X for given cover \mathcal{S} . \square

COROLLARY 4.8. *If X is a semi-Lindeloff space and each X -valued sequence has an $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}$ -cluster point, then X is a compact space.*

5. Some characterizations

Considering that semi-closed subsets of a semi-compact space is again semi-compact [13], we compare the collection of semi-limits of a sequence and that of semi-cluster points of a sequence in a topological space.

THEOREM 5.1. *Let \mathcal{I} and \mathcal{K} be two ideals satisfying ideality condition and K be a semi-compact subset of a space X . For any sequence $\{x_n\}$ in X , if $\{n \in \mathbb{N} : x_n \in K\} \notin (\mathcal{I} \cup \mathcal{K})$, then $K \cap sC_x(\mathcal{I}^{\mathcal{K}}) \neq \emptyset$.*

Proof. If possible, consider $K \cap sC_x(\mathcal{I}^{\mathcal{K}}) = \emptyset$. Then for each $c \in K$, there exists a set $M_c \in \mathcal{I}^*$ such that for any $U_c \in SO(X)$ containing c , we have $R_c = \{n \in M_c : x_n \in U_c\} \in \mathcal{K}$. Now, $K \subset \bigcup_{c \in K} U_c$, hence the semi open cover $\{U_c : c \in K\}$ contains a finite subcover $U_{c_1}, U_{c_2}, \dots, U_{c_k}$. Consider $M = M_1 \cap M_2 \cap \dots \cap M_k \in \mathcal{I}^*$ such that for each $c \in K$, we have $R_c = \{n \in M : x_n \in U_c\} \in \mathcal{K}$. So, $\{n \in M : x_n \in K\} \subset R_{c_1} \cup R_{c_2} \cup \dots \cup R_{c_k}$. Now, right hand side of the above set inequality belongs to \mathcal{K} , that implies $\{n \in M : x_n \in K\} \in \mathcal{K}$. But, $\{n \in \mathbb{N} : x_n \in K\} \subset \{n \in M : x_n \in K\} \cup M^c \in (\mathcal{I} \cup \mathcal{K})$. This is a contradiction. \square

THEOREM 5.2. *Let X be a semi-compact space. Suppose that $\{x_n\}$ is a sequence in X such that $sC_x(\mathcal{I}^{\mathcal{K}}) = \{l\}$. Then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}\text{-}\lim x_n = l$. Further, if the ideal \mathcal{K} is a P -ideal, then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}^*}\text{-}\lim x_n = l$.*

Proof. Consider a semi-open set U_l containing l . Then $X' = X \setminus U_l$ is a semi-closed subset of X . Now, for each $z \in X' \implies z \notin sC_x(\mathcal{I}^{\mathcal{K}})$. So, for all $M \in \mathcal{I}^*$, there exists $U_z \in SO(X)$ containing z such that $\{n \in M : x_n \in U_z\} \in \mathcal{K}$. Now, the semi-open cover $\{U_z\}_{z \in X'}$ of X' has a finite subcover, say $\{U_{z_1}, U_{z_2}, \dots, U_{z_k}\}$. Then $\bigcup_{i \leq k} \{n \in M : x_n \in U_{z_i}\} \in \mathcal{K}$. That implies $\{n \in M : x_n \in U_l\} \in \mathcal{K}^*$. Therefore, $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}\text{-}\lim x_n = l$. \square

The following result on the collection $sC_x(\mathcal{I}^{\mathcal{K}})$ and $sL(\mathcal{I}^{\mathcal{K}})$ (the collection of semi-limits of a sequence) summarise the Theorem 5.1 and Theorem 5.2.

THEOREM 5.3. *Suppose that $\{x_n\}$ is a sequence in X such that $\{n \in \mathbb{N} : x_n \in K\} \notin (\mathcal{I} \cup \mathcal{K})$, for any semi-compact $K \subset X$. Then $\mathcal{S}\text{-}\mathcal{I}^{\mathcal{K}}\text{-}\lim x_n = l$ if and only if $sC_x(\mathcal{I}^{\mathcal{K}}) = \{l\}$.*

In this segment, the existence problem mentioned in earlier sections is discussed further and the following results are obtained. Here we discuss the existence of the ideal \mathcal{J} for a given \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -clustered sequence such that their corresponding semi-cluster point sets coincides.

LEMMA 5.4. *Let \mathcal{I}, \mathcal{K} be two ideals on \mathbb{N} satisfying ideality condition. Let $x = x_n$ be a sequence in X . If $\mathcal{J} =$ ideal generated by $(\mathcal{K} \cup J)$, for any $J \in \mathcal{I}$. Then $a \in sC_x(\mathcal{J})$ implies $a \in sC_x(\mathcal{I}^{\mathcal{K}})$.*

Proof. Contrapositively, let $a \notin sC_x(\mathcal{I}^{\mathcal{K}})$. Then there exists at least an open set $V \in SO(X)$ containing a for which, for all $M \in \mathcal{I}^*$ we have $\{n \in M : x_n \in V\} \in \mathcal{K}$. Particularly, for $M_j = J^{\mathcal{C}} \in \mathcal{I}^*$, we have $\{n \in \mathbb{N} : x_n \in V\} \subseteq \{n \in M_j : x_n \in V\} \cup M_j^{\mathcal{C}}$. That implies $\{n \in \mathbb{N} : x_n \in V\} \in (\mathcal{K} \cup J)$. Hence, $a \notin sC_x(\mathcal{J})$. \square

THEOREM 5.5. *Let X be a Hausdorff space and $x = \{x_n\}$ be a sequence in X . Then there exists an ideal \mathcal{J} such that $a \in sC_x(\mathcal{J})$ if and only if $a \in sC_x(\mathcal{I}^{\mathcal{K}})$ provided $\mathcal{I} \cup \mathcal{K}$ is an ideal.*

Proof. Consider, $a \notin sC_x(\mathcal{J})$, where \mathcal{J} is the ideal generated by $\mathcal{K} \cup J$, for any $J \in \mathcal{I}$, provided $(\mathcal{I} \cup \mathcal{K})$ is an ideal. We claim that $a \notin sC_x(\mathcal{I}^{\mathcal{K}})$. So, there exists $U_a \in SO(X)$ containing a such that $\{n \in \mathbb{N} : x_n \in U_a\} \in \mathcal{J}$. Let $J = M^{\mathcal{C}}$. Thus,

$$\{n \in M : x_n \in U_a\} \subseteq \{n \in \mathbb{N} : x_n \in U_a\} \setminus M^{\mathcal{C}}.$$

Therefore, $\{n \in M : x_n \in U_a\} \subseteq (\mathcal{K} \cup J) \setminus J \in \mathcal{K}$, for any $M \in \mathcal{I}^*$. Thus $a \notin sC_x(\mathcal{I}^{\mathcal{K}})$. Converse part of the proof is immediate by Lemma 3.13. \square

For a non Hausdorff space, does there exist an ideal \mathcal{J} for a given \mathcal{S} - $\mathcal{I}^{\mathcal{K}}$ -clustered sequence such that their corresponding set of semi-cluster points coincides?

Here we characterize the collection of semi-cluster points of a sequence as a known subsets in a topological space. Recall that a subset D of a topological space X is said to be dense in X if for any open set U , $U \cap D \neq \emptyset$. In an arbitrary space, the notion of dense set for semi-open sets is equivalent to that of open sets.

THEOREM 5.6 ([12, Theorem 2.4]). *Let X be a space and $D \subset X$. Then D is dense in X if and only if $U \cap D \neq \emptyset$ for every $U \in SO(X)$.*

DEFINITION 5.7. Let X be an arbitrary space. We say that X is a semi-closed hereditarily separable space if every semi-closed subsets of X is separable.

THEOREM 5.8. *Let \mathcal{I}, \mathcal{K} be two ideals on \mathbb{N} and X be a space. Then*

(i) *For $x = \{x_n\}_{n \in \mathbb{N}}$, a sequence in X ; $sC_x(\mathcal{I}^{\mathcal{K}})$ is a semi-closed set.*

(ii) *If X is semi-closed hereditarily separable and there exists a disjoint sequence of sets $\{D_n\}$ such that $D_n \subset \mathbb{N}$, $D_n \notin \mathcal{I}, \mathcal{K}$ for all n , then for every non empty semi-closed subset F of X , there exists a sequence x in X such that $F = sC_x(\mathcal{I}^{\mathcal{K}})$ provided $\mathcal{I} \cup \mathcal{K}$ is an ideal.*

Proof. Consider the sequence $x = \{x_n\}$ in X and let \mathcal{I}, \mathcal{K} be the two ideals on \mathbb{N} .

(i) Let $y \in sCl(C_x(\mathcal{I}^{\mathcal{K}}))$; the semi-closure of $C_x(\mathcal{I}^{\mathcal{K}})$. Let U be a semi-open set containing y . It is clear that $U \cap C_x(\mathcal{I}^{\mathcal{K}}) \neq \emptyset$. Let $q \in U \cap C_x(\mathcal{I}^{\mathcal{K}})$ i.e., $q \in U$ and $q \in C_x(\mathcal{I}^{\mathcal{K}})$. Now there exists a set $M \in \mathcal{I}^*$ such that $\{n \in M : y_n \in U\} \notin \mathcal{K}$. Thus, $y \in C_x(\mathcal{I}^{\mathcal{K}})$.

(ii) F is separable as a semi-closed subset of X . Then by Definition 2.3 of [15] and Theorem 5.6, there exists a countable set $S = \{s_1, s_2, \dots\} \subset F$ such that $sCl(S) = F$. Consider $x_n = s_i$ for $n \in D_i$. Then, we have a subsequence $\{k_n\}$ of the sequence $\{n\}$. Now, consider the sequence $x = \{x_{n_k}\}$ and let $y \in sC_x(\mathcal{K})$ (taking $y \neq s_i$ otherwise if $y = s_i$ for some i , then y is eventually (except finite elements) in F). We claim that $sC_x(\mathcal{I}^{\mathcal{K}}) \subset F$. Let U be any semi-open set containing y . Then $\{n : x_{n_k} \in U\} \notin \mathcal{K}$ ($\neq \emptyset$). So, $s_i \in U$ for some i . Therefore, $F \cap U$ is non empty. So y is a semi-limit point of F and semi-closedness of F implies $y \in F$. Hence $sC_x(\mathcal{K}) \subset F$. Further $sC_x(\mathcal{I}^{\mathcal{K}}) \subseteq sC_x(\mathcal{K}) \subset F$. For the converse argument, consider $a \in F$ and U be a semi-open set containing a , then there exists $s_i \in S$ such that $s_i \in U$. Then $\{n : x_{n_k} \in U\} \supset D_i$ ($\notin \mathcal{K}, \mathcal{I}$). Thus $\{n : x_{n_k} \in U\} \notin (\mathcal{I} \cup \mathcal{K})$ i.e., $a \in sC_x(\mathcal{I} \cup \mathcal{K})$. Again, by Lemma 4.3, $sC_f(\mathcal{I} \cup \mathcal{K}) \subseteq sC_f(\mathcal{I}^{\mathcal{K}})$. Thus, we get the reverse implication. \square

REMARK 5.9. Theorem 5.8 extends [8, Theorem 10] to semi-open sets, it follows by letting \mathcal{I}, \mathcal{K} coincide and using open sets instead of semi-open sets in the above theorem.

ACKNOWLEDGEMENT. The authors would like to thank the referees for their insightful comments that helped to enhance the clarity and presentation of the article. The first author would also like to thank the University Grants Commission (UGC) for awarding the JRF vide UGC-Ref. No.: 1115/(CSIR-UGC NET DEC. 2017), India.

REFERENCES

- [1] S. G. Crossley, S. K. Hildebrand, *Semi-closure*. Texas J. Sci., **22(2-3)** (1970), 99–112.
- [2] S. G. Crossley, S. K. Hildebrand, *Semi-topological properties*, Fund. Math., **74(3)** (1972), 233–254.
- [3] P. Das, S. Sengupta, J. Supina, *$\mathcal{I}^{\mathcal{K}}$ -convergence of sequence of functions*, Math. Slovaca, **69(5)**(2019), 1137–1148.
- [4] P. Das, S. Dasgupta, S. Glab, M. Bienias, *Certain aspects of ideal convergence in topological spaces*, Topology Appl., **275** (2020), Article 107005.
- [5] A. Guevara, J. Sanabria, E. Rosas, *\mathcal{S} - \mathcal{I} -convergence of sequences*, Trans. A. Razmadze Math. Inst., **174** (2020), 75–81.
- [6] K. P. Hart, J. Nagata, J. E. Vaughan. *Encyclopedia of General Topology*, Elsevier Science Publications, Amsterdam-Netherlands, 2004.
- [7] P. Kostyrko, T. Salat, W. Wilczynski, *\mathcal{I} -convergence*, Real Anal. Exchange, **26** (2001), 669–685.
- [8] B. K. Lahiri, P. Das, *\mathcal{I} and \mathcal{I}^* -convergence in topological spaces*, Math. Bohem., **130** (2005), 153–160.
- [9] N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, **70** (1963), 36–41.
- [10] M. Macaj, M. Slezia, *$\mathcal{I}^{\mathcal{K}}$ -convergence*, Real Anal. Exchange, **36** (2011), 177–194.

- [11] S. N. Maheshwari, R. Prasad, *Some new separation axioms*, Ann. Soc. Sci. Bruxelles Ser. I, **89(3)**(1975), 395–402.
- [12] S. Modak, *Remarks on dense set*, Int. Math. Forum, **6** (2011), 2153–2158.
- [13] M. S. Sarsak, *On semicompact sets and associated properties*, Int. J. Math. Math. Sci., **2009** (2009), Article ID 465387.
- [14] A. Sharmah, D. Hazarika, *Further aspects of $\mathcal{I}^{\mathcal{K}}$ -convergence in topological spaces*, Appl. Gen. Topol., **22(2)** (2021), 355–366.
- [15] A. Sharmah, D. Hazarika, *Some properties of $\mathcal{I}^{\mathcal{K}}$ -convergence*, J. Math. Comput. Sci., **11(2)** (2021), 1528–1534.

(received 07.12.2022; in revised form 26.03.2024; available online 19.09.2024)

Department of Mathematics, Bhattadev University, Pathsala 781325, Assam, India

E-mail: ankursarmah189@gmail.com

ORCID iD: <https://orcid.org/0000-0002-3894-642X>

Department of Mathematical Sciences, Tezpur University, Napaam 784028, Assam, India

E-mail: debajit@tezu.ernet.in

ORCID iD: <https://orcid.org/0000-0002-1145-3721>