

FIBONACCI NUMBERS WHICH ARE CONCATENATIONS OF THREE REPDIGITS

Fatih Erduvan and Refik Keskin

Abstract. In this study, it is proved that the only Fibonacci numbers which are concatenations of three repdigits are 144, 233, 377, 610, 987, 17711.

1. Introduction

Let (F_n) be the sequence of Fibonacci numbers given by $F_0 = 0$, $F_1 = 1$, and $F_n = F_{n-1} + F_{n-2}$, for $n \geq 2$. Binet formula for the n^{th} Fibonacci number is

$$F_n = \frac{\alpha^n - \beta^n}{\sqrt{5}},$$

where $\alpha = \frac{1 + \sqrt{5}}{2}$ and $\beta = \frac{1 - \sqrt{5}}{2}$ are the roots of the characteristic equation $x^2 - x - 1 = 0$. It can be seen that $1 < \alpha < 2$, $-1 < \beta < 0$ and $\alpha\beta = -1$. A relation between n^{th} Fibonacci number F_n and α is given by

$$\alpha^{n-2} \leq F_n \leq \alpha^{n-1} \quad \text{for } n \geq 1. \quad (1)$$

The inequality (1) can be proved by induction. Given $k \geq 1$, we say that N is a concatenations of k repdigits, if N can be written in the form

$$\overbrace{d_1 \dots d_1}^{m_1 \text{ times}} \overbrace{d_2 \dots d_2}^{m_2 \text{ times}} \dots \overbrace{d_k \dots d_k}^{m_k \text{ times}}.$$

In [1], the authors solved the problem of finding the Fibonacci numbers which are concatenations of two repdigits. In [6, 7] Ddamulira determined all the tribonacci numbers that are concatenations of two repdigits, respectively. In [10], Trojovský considered Fibonacci numbers of the form

$$F_n = a \overbrace{b \dots b}^{m \text{ times}} \overbrace{c \dots c}^{k \text{ times}},$$

2020 Mathematics Subject Classification: 11B39, 11J86, 11D61

Keywords and phrases: Fibonacci number; concatenations; repdigit; Diophantine equations; linear forms in logarithms.

where $a \geq 1, 0 \leq a, b, c \leq 9$ with $2 \leq k \leq m$. He showed that the largest Fibonacci numbers of the above form is 17711. Motivated by these studies, in this paper, we study the equation

$$F_n = \underbrace{\overline{d_1 \dots d_1}}_{m_1 \text{ times}} \underbrace{\overline{d_2 \dots d_2}}_{m_2 \text{ times}} \underbrace{\overline{d_3 \dots d_3}}_{m_3 \text{ times}}, \tag{2}$$

where $d_1, m_1, m_2, m_3 \geq 1$ and $0 \leq d_1, d_2, d_3 \leq 9$. That is, we find all Fibonacci numbers which are concatenations of three repdigits. It is shown that the only Fibonacci numbers which are concatenations of three repdigits are 144, 233, 377, 610, 987, 17711. In Section 2, we introduce necessary lemmas. Then we prove our main theorem in Section 3.

2. Auxiliary results

Let η be an algebraic number of degree d with minimal polynomial

$$a_0x^d + a_1x^{d-1} + \dots + a_d = a_0 \prod_{i=1}^d (x - \eta^{(i)}) \in \mathbb{Z}[x],$$

where the a_i 's are relatively prime integers with $a_0 > 0$ and the $\eta^{(i)}$'s are conjugates of η . Then

$$h(\eta) = \frac{1}{d} \left(\log a_0 + \sum_{i=1}^d \log \left(\max \{ |\eta^{(i)}|, 1 \} \right) \right) \tag{3}$$

is called the logarithmic height of η . In particular, if $\eta = a/b$ is a rational number with $\gcd(a, b) = 1$ and $b \geq 1$, then $h(\eta) = \log(\max\{|a|, b\})$.

For algebraic numbers η and γ , the function h has the following basic properties (see [4]):

$$h(\eta \pm \gamma) \leq h(\eta) + h(\gamma) + \log 2, \quad h(\eta\gamma^{\pm 1}) \leq h(\eta) + h(\gamma), \quad h(\eta^m) = |m|h(\eta).$$

Now, we give a theorem which is deduced from [9, Corollary 2.3] and provides a large upper bound for the subscript n in the equations (2) (also see [5, Theorem 9.4]).

LEMMA 2.1. *Assume that $\gamma_1, \gamma_2, \dots, \gamma_t$ are positive real algebraic numbers in a real algebraic number field \mathbb{K} of degree D , b_1, b_2, \dots, b_t are rational integers, and $\Lambda := \gamma_1^{b_1} \dots \gamma_t^{b_t} - 1$ is not zero. Then*

$$|\Lambda| > \exp(-1.4 \cdot 30^{t+3} \cdot t^{4.5} \cdot D^2(1 + \log D)(1 + \log B)A_1A_2 \dots A_t),$$

where $B \geq \max\{|b_1|, \dots, |b_t|\}$ and $A_i \geq \max\{Dh(\gamma_i), |\log \gamma_i|, 0.16\}$ for all $i = 1, \dots, t$.

The following lemma is given in [3]. This lemma is an immediate variation of the result due to Dujella and Pethő from [8], which is a version of a lemma of Baker and Davenport [2]. It will be used to reduce the upper bound for the subscript n in the equation (2). Let $\|x\| = \min\{|x - n| : n \in \mathbb{Z}\}$ for any real number x .

LEMMA 2.2. *Let M be a positive integer, let p/q be a convergent of the continued fraction of the irrational number γ such that $q > 6M$, and let A, B, μ be some real*

numbers with $A > 0$ and $B > 1$. Let $\epsilon := \|\mu q\| - M\|\gamma q\|$. If $\epsilon > 0$, then there exists no solution to the inequality $0 < |u\gamma - v + \mu| < AB^{-w}$, in positive integers u, v , and w with $u \leq M$ and $w \geq \frac{\log(Aq/\epsilon)}{\log B}$.

The following lemma is given in [11].

LEMMA 2.3. Let $a, x \in \mathbb{R}$. If $0 < a < 1$ and $|x| < a$; then

$$|\log(1+x)| < \frac{-\log(1-a)}{a} \cdot |x| \quad \text{and} \quad |x| < \frac{a}{1-e^{-a}} \cdot |e^x - 1|.$$

3. Main theorem

THEOREM 3.1. The only Fibonacci numbers which are concatenations of three repdigits are 144, 233, 377, 610, 987, 17711.

Proof. Assume that the equation (2) holds. Since the number F_{149} has 31 digits, $1 \leq m_1, m_2, m_3 \leq 29$ for $n \leq 149$. Then by using Mathematica program, we can find all the solutions to the equation (2) for $d_1 > 0$, $0 \leq d_1, d_2, d_3 \leq 9$, and $1 \leq n \leq 149$. In this case, we see that $F_n \in \{144, 233, 377, 610, 987, 17711\}$. From now on, we assume that $n \geq 150$ and we rule out $d_1 = d_2 \neq d_3$ and $d_1 \neq d_2 = d_3$ in equation (2) since these cases have been solved in [1]. Furthermore, the case $d_1 = d_2 = d_3$ in equation (2) is impossible since the largest repdigit in Fibonacci sequence is $F_{10} = 55$, which has two digits. As

$$F_n = \underbrace{\overbrace{d_1 \dots d_1}_{m_1 \text{ times}} \overbrace{d_2 \dots d_2}_{m_2 \text{ times}} \overbrace{d_3 \dots d_3}_{m_3 \text{ times}}}_{m_1 \text{ times } m_2 \text{ times } m_3 \text{ times}} = \underbrace{d_1 \dots d_1}_{m_1 \text{ times}} \times 10^{m_2+m_3} + \underbrace{d_2 \dots d_2}_{m_2 \text{ times}} \times 10^{m_3} + \underbrace{d_3 \dots d_3}_{m_3 \text{ times}}$$

we get
$$F_n = \frac{d_1(10^{m_1} - 1)}{9} 10^{m_2+m_3} + \frac{d_2(10^{m_2} - 1)}{9} 10^{m_3} + \frac{d_3(10^{m_3} - 1)}{9}, \tag{4}$$

i.e.
$$F_n = \frac{1}{9} (d_1 10^{m_1+m_2+m_3} - (d_1 - d_2) 10^{m_2+m_3} - (d_2 - d_3) 10^{m_3} - d_3). \tag{5}$$

Combining the right-hand side of inequality of (1) with (4), we obtain

$$10^{m_1+m_2+m_3-1} < F_n \leq \alpha^{n-1} < 10^{n-1}.$$

From this, we get $m_1 + m_2 + m_3 < n$. Now, we can rewrite equation (5) as

$$\begin{aligned} & 9\alpha^n - d_1\sqrt{5} \cdot 10^{m_1+m_2+m_3} \\ & = 9\beta^n - (d_1 - d_2)\sqrt{5} \cdot 10^{m_2+m_3} - (d_2 - d_3)\sqrt{5} \cdot 10^{m_3} - d_3\sqrt{5}. \end{aligned} \tag{6}$$

Taking absolute values of both sides of the equation (6), we get

$$\begin{aligned} & \left| 9\alpha^n - d_1\sqrt{5} \cdot 10^{m_1+m_2+m_3} \right| \\ & \leq 9|\beta|^n + |d_1 - d_2|\sqrt{5} \cdot 10^{m_2+m_3} + |d_2 - d_3|\sqrt{5} \cdot 10^{m_3} + d_3\sqrt{5} \\ & \leq 9\alpha^{-n} + 9\sqrt{5} \cdot 10^{m_2+m_3} + 9\sqrt{5} \cdot 10^{m_3} + 9\sqrt{5} \\ & \leq 9\alpha^{-n} + \sqrt{5}(9 \cdot 10^{m_2+m_3} + 9 \cdot 10^{m_3} + (0.9) \cdot 10^{m_3}) \end{aligned}$$

$$= 9\alpha^{-n} + \sqrt{5} \cdot 10^{m_3}(9 \cdot 10^{m_2} + (9.9)) \leq 9\alpha^{-n} + \sqrt{5} \cdot 10^{m_3}(9 \cdot 10^{m_2} + (0.99) \cdot 10^{m_2}) \\ < (0.09)\alpha^{-n}10^{m_2+m_3} + 22.34 \cdot 10^{m_2+m_3} < 22.341 \cdot 10^{m_2+m_3},$$

where we have used the fact that $n \geq 150$. Therefore, it is seen that

$$\left| 9\alpha^n - d_1\sqrt{5} \cdot 10^{m_1+m_2+m_3} \right| < 22.341 \cdot 10^{m_2+m_3}. \quad (7)$$

Dividing both sides of (7) by $d_1\sqrt{5} \cdot 10^{m_1+m_2+m_3}$, we obtain

$$\left| \frac{9}{d_1\sqrt{5}}\alpha^n 10^{-m_1-m_2-m_3} - 1 \right| \leq \frac{22.341 \cdot 10^{m_2+m_3}}{d_1\sqrt{5} \cdot 10^{m_1+m_2+m_3}} < \frac{9.992}{10^{m_1}}. \quad (8)$$

Now, let us apply Lemma 2.1 with $\gamma_1 := \frac{9}{d_1\sqrt{5}}, \gamma_2 := \alpha, \gamma_3 := 10$ and $b_1 := 1, b_2 := n, b_3 := -m_1 - m_2 - m_3$. Note that the numbers γ_1, γ_2 , and γ_3 are positive real numbers and elements of the field $K = \mathbb{Q}(\sqrt{5})$. The degree of the field K is 2. So $D = 2$. It can be seen that $\Lambda_1 := \frac{9}{d_1\sqrt{5}}\alpha^n 10^{-m_1-m_2-m_3} - 1$ is nonzero. Moreover, since

$$h(\gamma_1) = h\left(\frac{9}{d_1\sqrt{5}}\right) \leq h\left(\frac{9}{d_1}\right) + h(\sqrt{5}) \leq \log 9 + \frac{\log 5}{2} < 3.1,$$

and
$$h(\gamma_2) = h(\alpha) = \frac{\log \alpha}{2} < 0.25, \quad h(\gamma_3) = h(10) = \log 10 < 2.31$$

by (3), we can take $A_1 := 6.2, A_2 := 0.5$, and $A_3 := 4.62$. On the other hand, as $m_1 + m_2 + m_3 < n$ and $B \geq \max\{|1|, |n|, |-m_1 - m_2 - m_3|\}$, we can take $B := n$. Thus, taking into account the inequality (8) and using Lemma 2.1, we obtain

$$(9.992) \cdot 10^{-m_1} > |\Lambda_1| > \exp(C \cdot (1 + \log n)),$$

where $C = -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2 \cdot (1 + \log 2) \cdot (6.2) \cdot (0.5) \cdot (4.62)$. By a simple computation, it follows that

$$m_1 \log 10 < 1.39 \cdot 10^{13} \cdot (1 + \log n) + \log(9.992). \quad (9)$$

Rearranging the equation (5) as

$$9\alpha^n - d_1\sqrt{5} \cdot 10^{m_1+m_2+m_3} \\ = 9\beta^n - (d_1 - d_2)\sqrt{5} \cdot 10^{m_2+m_3} - (d_2 - d_3)\sqrt{5}10^{m_3} - d_3\sqrt{5} \quad (10)$$

and taking absolute values of both sides of the equation (10), we get

$$\left| 9\alpha^n - (d_1 10^{m_1} - (d_1 - d_2))\sqrt{5} \cdot 10^{m_2+m_3} \right| \\ \leq 9|\beta|^n + |d_2 - d_3|\sqrt{5} \cdot 10^{m_3} + d_3\sqrt{5} \leq 9\alpha^{-n} + 9\sqrt{5} \cdot 10^{m_3} + 9\sqrt{5} \\ \leq 9\alpha^{-n} + \sqrt{5}(9 \cdot 10^{m_3} + 9) \leq 9\alpha^{-n} + \sqrt{5}(9 \cdot 10^{m_3} + (0.9) \cdot 10^{m_3}) < 22.15 \cdot 10^{m_3},$$

i.e.,
$$\left| 9\alpha^n - (d_1 10^{m_1} - (d_1 - d_2))\sqrt{5} \cdot 10^{m_2+m_3} \right| < 22.15 \cdot 10^{m_3}. \quad (11)$$

Dividing both sides of (11) by $(d_1 10^{m_1} - (d_1 - d_2))\sqrt{5} \cdot 10^{m_2+m_3}$, we obtain

$$\left| 1 - \left(\frac{9}{(d_1 10^{m_1} - (d_1 - d_2))\sqrt{5}} \right) \alpha^n 10^{-m_2-m_3} \right| < \frac{1.11}{10^{m_2}}. \quad (12)$$

Taking $\gamma_1 := \frac{9}{(d_1 10^{m_1} - (d_1 - d_2))\sqrt{5}}, \gamma_2 := \alpha, \gamma_3 := 10$ and $b_1 := 1, b_2 := n, b_3 :=$

$-m_2 - m_3$, we can apply Lemma 2.1. The numbers γ_1, γ_2 , and γ_3 are positive real numbers and elements of the field $K = \mathbb{Q}(\sqrt{5})$ and so $D = 2$. One can verify that $\Lambda_2 := 1 - \left(\frac{9}{(d_1 10^{m_1} - (d_1 - d_2))\sqrt{5}}\right) \alpha^n 10^{-m_2 - m_3}$ is nonzero. By using (3) and the properties of the logarithmic height, we get

$$\begin{aligned} h(\gamma_1) &= h\left(\frac{9}{(d_1 10^{m_1} - (d_1 - d_2))\sqrt{5}}\right) \\ &\leq h(9) + h(\sqrt{5}) + h(d_1 10^{m_1}) + h(d_1 - d_2) + \log 2 \\ &\leq 3 \log 9 + \frac{\log 5}{2} + m_1 \log 10 + \log 2 < 8.1 + m_1 \log 10, \\ h(\gamma_2) &= h(\alpha) = \frac{\log \alpha}{2} < 0.25, \end{aligned}$$

and $h(\gamma_3) = h(10) = \log 10 < 2.31$.

So, we can take $A_1 := 16.2 + 2m_1 \log 10$, $A_2 := 0.5$, and $A_3 := 4.62$. As $m_2 + m_3 < n$ and $B \geq \max\{|1|, | -n|, | -m_2 - m_3|\}$, we can take $B := n$. Thus, taking into account the inequality (12) and using Lemma 2.1, we obtain

$$1.11 \cdot 10^{-m_2} > |\Lambda_2| > \exp(C \cdot (1 + \log n) (16.2 + 2m_1 \log 10))$$

i.e., $m_2 \log 10 < 2.25 \cdot 10^{12} \cdot (1 + \log n) (16.2 + 2m_1 \log 10) + \log(1.11)$, (13)

where $C = -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2 (1 + \log 2) \cdot (0.5) \cdot (4.62)$. Rearranging the equation (5) as

$$9\alpha^n - (d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5} \cdot 10^{m_3} = 9\beta^n - d_3$$
 (14)

and taking absolute values of both sides of the equation (14), we get

$$\left| 9\alpha^n - (d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5} \cdot 10^{m_3} \right| \leq 9|\beta|^n + d_3 = 9\alpha^{-n} + 9 < 9.1,$$

i.e., $\left| 9\alpha^n - (d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5} \cdot 10^{m_3} \right| < 9.1$. (15)

If we divide both sides of (15) by $9\alpha^n$, we obtain

$$\left| 1 - \left(\frac{(d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}}{9}\right) \alpha^{-n} 10^{m_3} \right| \leq 1.02 \cdot \alpha^{-n}. \quad (16)$$

Taking $\gamma_1 := \left(\frac{(d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}}{9}\right)$, $\gamma_2 := \alpha$, $\gamma_3 := 10$ and $b_1 := 1$, $b_2 := -n$, $b_3 := m_3$, we can apply Lemma 2.1. The numbers γ_1, γ_2 and γ_3 are positive real numbers and elements of the field $K = \mathbb{Q}(\sqrt{5})$ and so $D = 2$. One can verify that

$$\Lambda_3 := 1 - \left(\frac{(d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}}{9}\right) \alpha^{-n} 10^{m_3}$$

is nonzero. By using (3) and the properties of the logarithmic height, we get

$$\begin{aligned} h(\gamma_1) &= h\left(\frac{(d_1 10^{m_1 + m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}}{9}\right) \\ &\leq h(\sqrt{5}) + h(9) + h(d_1 10^{m_1 + m_2}) + h((d_1 - d_2) 10^{m_2}) + h(d_2 - d_3) + 2 \log 2 \end{aligned}$$

$$\begin{aligned} &\leq \frac{\log 5}{2} + 4 \log 9 + (m_1 + m_2) \log 10 + m_2 \log 10 + 2 \log 2 \\ &< 10.98 + (m_1 + m_2) \log 10 + m_2 \log 10, \end{aligned}$$

$$h(\gamma_2) = h(\alpha) = \frac{\log \alpha}{2} < 0.25, \quad \text{and} \quad h(\gamma_3) = h(10) = \log 10 < 2.31.$$

So, we can take $A_1 := 21.96 + 2m_1 \log 10 + 4m_2 \log 10$, $A_2 := 0.5$, and $A_3 := 4.62$. As $m_3 < n$ and $B \geq \max\{|1|, |-n|, |m_3|\}$, we can take $B := n$. Thus, taking into account the inequality (16) and using Lemma 2.1, we obtain

$$\begin{aligned} &1.02 \cdot \alpha^{-n} > |\Lambda_3| > \exp(C \cdot (1 + \log n) (21.96 + 2m_1 \log 10 + 4m_2 \log 10)) \\ \text{i.e.,} \quad &n \log \alpha - \log(1.02) < 2.25 \cdot 10^{12} (1 + \log n) (21.96 + 2m_1 \log 10 + 4m_2 \log 10), \end{aligned} \quad (17)$$

where $C = -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2 (1 + \log 2) \cdot (0.5) \cdot (4.62)$. Using the inequalities (9), (13) and (17), a computer search with Mathematica gives us that $n < 1.36 \cdot 10^{45}$.

Now, let us try to reduce the upper bound on n by applying Lemma 2.2. Let

$$z_1 := (m_1 + m_2 + m_3) \log 10 - n \log \alpha - \log \left(\frac{9}{\sqrt{5}d_1} \right).$$

From (8), it is seen that $|x| = |e^{-z_1} - 1| < \frac{9.992}{10^{m_1}} < 0.9995$ for $m_1 \geq 1$. Choosing $a := 0.9995$, by Lemma 2.3, we get the inequality

$$|z_1| = |\log(x + 1)| < \frac{\log(2000)}{0.9995} \cdot \frac{9.992}{10^{m_1}} < \frac{75.99}{10^{m_1}}.$$

Thus, it follows that

$$0 < \left| (m_1 + m_2 + m_3) \log 10 - n \log \alpha - \log \left(\frac{9}{\sqrt{5}d_1} \right) \right| < \frac{75.99}{10^{m_1}}.$$

Dividing this inequality by $\log \alpha$, we get

$$0 < \left| (m_1 + m_2 + m_3) \frac{\log 10}{\log \alpha} - n - \left(\frac{\log(9/\sqrt{5}d_1)}{\log \alpha} \right) \right| < 157.92 \cdot 10^{-m_1}. \quad (18)$$

Now, we can apply Lemma 2.2. Put $\gamma := \frac{\log 10}{\log \alpha} \notin \mathbb{Q}$, $\mu := -\frac{\log(9/\sqrt{5}d_1)}{\log \alpha}$, $A := 157.92$, $B := 10$ and $w := m_1$. Let $M := 1.36 \cdot 10^{45}$. Then $M > m_1 + m_2 + m_3$ and denominator of the 97th convergent of γ exceeds $6M$. Furthermore, $\epsilon := \|\mu q_{97}\| - M \|\gamma q_{97}\| > 0.12$.

Thus, the inequality (18) has no solutions for $\frac{\log(Aq_{97}/\epsilon)}{\log B} < 49.62 < m_1$. So $m_1 \leq 49$.

Now replace (13) by (17). Substituting this upper bound for m_1 into (17), we obtain $n < 1.12 \cdot 10^{32}$. Now, let

$$z_2 := (m_2 + m_3) \log 10 - n \log \alpha - \log \left(\frac{9}{(d_1 10^{m_1} - (d_1 - d_2)) \sqrt{5}} \right).$$

From (12), it is seen that $|x| = |e^{-z_2} - 1| < 1.11 \cdot 10^{-m_2} < 0.2$ for $m_2 \geq 1$. Choosing $a := 0.2$, we get the inequality

$$|z_2| = |\log(x + 1)| < \frac{\log(10/8)}{0.2} \cdot \frac{1.11}{10^{m_2}} < \frac{1.24}{10^{m_2}}$$

by Lemma 2.3. Thus, it follows that

$$0 < \left| (m_2 + m_3) \log 10 - n \log \alpha - \log \left(\frac{9}{(d_1 10^{m_1} - (d_1 - d_2)) \sqrt{5}} \right) \right| < 1.24 \cdot 10^{-m_2}.$$

Dividing both sides of the above inequality by $\log \alpha$, shows that

$$0 < \left| (m_2 + m_3) \frac{\log 10}{\log \alpha} - n + \frac{\log (9/(d_1 10^{m_1} - (d_1 - d_2)) \sqrt{5})}{\log \alpha} \right| < 2.57 \cdot 10^{-m_2}. \quad (19)$$

Putting $\gamma := \frac{\log 10}{\log \alpha}$ and taking $m_2 + m_3 < M := 5.57 \cdot 10^{31}$, we find that q_{73} , the denominator of the 73rd convergent of γ exceeds $6M$. Taking $\mu := \frac{\log(9/(d_1 10^{m_1} - (d_1 - d_2)) \sqrt{5})}{\log \alpha}$ and considering the fact that $m_1 \leq 49$, $d_1 \neq d_2$, $1 \leq d_1 \leq 9$ and $0 \leq d_2 \leq 9$, a quick computation with Mathematica gives us the equality $\epsilon = \epsilon(\mu) := \|\mu q_{73}\| - M \|\gamma q_{73}\| > 0.00001$. Let $A := 2.57$, $B := 10$ and $w := m_2$ in Lemma 2.2. Then with the help of Mathematica, we can say that the inequality (19) has no solution for $\frac{\log(Aq_{73}/\epsilon)}{\log B} < 39.51 < m_2$. Therefore $m_2 \leq 42$. Substituting this upper bound for m_1 and m_2 into (17), we obtain $n < 1.15 \cdot 10^{17}$. Now, let

$$z_3 := m_3 \log 10 - n \log \alpha + \log \left(\frac{(d_1 10^{m_1+m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}}{9} \right).$$

From (16), one can write $|x| = |e^{z_3} - 1| < 1.02 \cdot \alpha^{-n} < 0.01$ for $n \geq 150$. Choosing $a := 0.01$, we get the inequality

$$|z_3| = |\log(x + 1)| < \frac{\log(100/99)}{0.01} \cdot \frac{1.02}{\alpha^n} < \frac{1.03}{\alpha^n}$$

by Lemma 2.3. Thus, it follows that

$$0 < \left| m_3 \log 10 - n \log \alpha + \log \left(\frac{d_1 10^{m_1+m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)}{9} \right) \right| < \frac{1.03}{\alpha^n}.$$

Dividing both sides of the above inequality by $\log \alpha$, we get

$$0 < \left| \frac{m_3 \log 10}{\log \alpha} - n + \frac{\log ((d_1 10^{m_1+m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}/9)}{\log \alpha} \right| < \frac{2.15}{\alpha^n}. \quad (20)$$

Putting $\gamma := \frac{\log 10}{\log \alpha}$ and taking $M := 1.15 \cdot 10^{17}$, it is seen that $M > m_3$ and q_{48} , the denominator of the 48th convergent of γ exceeds $6M$.

Taking $\mu := \frac{\log((d_1 10^{m_1+m_2} - (d_1 - d_2) 10^{m_2} - (d_2 - d_3)) \sqrt{5}/9)}{\log \alpha}$ and considering the fact that $m_1 \leq 49$, $m_2 \leq 39$, $1 \leq d_1 \leq 9$ and $0 \leq d_2, d_3 \leq 9$, a quick computation with Mathematica gives us the equality $\epsilon = \epsilon(\mu) := \|\mu q_{48}\| - M \|\gamma q_{48}\| > 0$ except for the case $d_1 = d_2 \neq d_3$, $d_1 \neq d_2 = d_3$ and $d_1 = d_2 = d_3$. Let $A := 2.15$, $B := \alpha$ and $w := n$ in Lemma 2.2. Then with the help of Mathematica, we can say that the inequality (20) has no solution for $\frac{\log(Aq_{48}/\epsilon)}{\log B} < 147.17 < n$. Therefore $n \leq 147$. This contradicts our assumption that $n \geq 150$. This completes the proof. \square

REFERENCES

- [1] A. Alahmadi, A. Altassan, F. Luca, H. Shoaib, *Fibonacci numbers which are concatenations of two repdigits*, Quaest. Math., **44(2)** (2021), 281–290.
- [2] A. Baker and H. Davenport, *The equations $3x^2 - 2 = y^2$ and $8x^2 - 7 = z^2$* , Quart. J. Math. Oxford Ser. (2), **20(1)** (1969), 129–137.
- [3] J.J. Bravo, C.A. Gomez, F. Luca, *Powers of two as sums of two k -Fibonacci numbers*, Miskolc Math. Notes, **17(1)** (2016), 85–100.
- [4] Y. Bugeaud, *Linear Forms in Logarithms and Applications*, IRMA Lectures in Mathematics and Theoretical Physics, 28, Zurich: European Mathematical Society, 2018.
- [5] Y. Bugeaud, M. Mignotte, S. Siksek, *Classical and modular approaches to exponential Diophantine equations I. Fibonacci and Lucas perfect powers*, Ann. of Math. **163(3)** (2006), 969–1018.
- [6] M. Ddamulira, *Tribonacci numbers that are concatenations of two repdigits*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM, **114(4)** (2020), 203.
- [7] M. Ddamulira, *Padovan numbers that are concatenations of two repdigits*, Math. Slovaca, **71(2)** (2021), 275–284.
- [8] A. Dujella, A. Pethò, *A generalization of a theorem of Baker and Davenport*, Quart. J. Math. Oxford Ser. (2), **49(3)** (1998), 291–306.
- [9] E. M. Matveev, *An explicit lower bound for a homogeneous rational linear form in the logarithms of algebraic numbers II*, Izv. Ross. Akad. Nauk Ser. Mat., **64(6)** (2000), 125–180 (Russian). Translation in Izv. Math. **64(6)** (2000), 1217–1269.
- [10] P. Trojovský, *On Fibonacci numbers with a prescribed block of digits*, Mathematics **8(4)** (2020), 639.
- [11] B. M. M. de Weger, *Algorithms for Diophantine Equations*, CWI Tracts 65, Stichting Maths. Centrum, Amsterdam, 1989.

(received 22.08.2020; in revised form 29.05.2021; available online 31.01.2022)

Sakarya University, Mathematics Department, Sakarya, Turkey

E-mail: erduvanmat@hotmail.com

Sakarya University, Mathematics Department, Sakarya, Turkey

E-mail: rkeskin@sakarya.edu.tr