

CONVERGENCE OF FINITE-DIFFERENCE SCHEMES FOR POISSON'S EQUATION WITH BOUNDARY CONDITION OF THE THIRD KIND

Branislav Popović

Abstract. In this paper we study the convergence of finite-difference schemes to generalized solutions of the third boundary-value problem for Poisson's equation on the unit square. Using the generalized Bramble-Hilbert lemma, we obtain error estimates in discrete H^1 Sobolev norm compatible, in some cases, with the smoothness of the data.

The outline of the paper is as follows. In section 1 notational conventions are presented. The stability theorem is proved in section 2. In section 3 we prove estimates of the energy of the operator Δ_h . Finally, in section 4, we derive our main results.

1. Preliminaries and notation

Consider the third boundary-value problem for Poisson's equation on the unit square $\Omega = (0, 1)^2$:

$$\Delta u = f \quad \text{in } \Omega, \quad \frac{\partial u}{\partial n} + \sigma u = 0 \quad \text{on } \partial\Omega \quad (1)$$

where $\frac{\partial u}{\partial n} = \frac{\partial u}{\partial x} \cos(x, n) + \frac{\partial u}{\partial y} \cos(y, n)$, n the unit outward normal to $\partial\Omega$, and $\sigma(x, y)$ is continuous function on $\partial\Omega$ such that $\sigma(x, y) \geq \sigma_0 > 0$, $\sigma_0 = \text{const}$.

We suppose that, for $f \in H^0(\Omega)$, our problem (1) has a unique solution in $H^2(\Omega)$ and, provided $f \in H^{s-2}(\Omega)$, $u \in H^s(\Omega)$ for $2 \leq s \leq 4$ (see [1,4]).

Problem (1) is discretised on the uniform mesh with step-size $h : \bar{\Omega}_h = \{(ih, jh) : i, j = 0, 1, 2, \dots, N; Nh = 1\}$. We define $\Omega_h = \Omega \cap \bar{\Omega}_h$ and $\partial\Omega_h = \partial\Omega \cap \bar{\Omega}_h$. In $\partial\Omega_h$ we distinguish between two kinds of meshpoints: $\partial\Omega_h^2 = \partial\Omega_h \setminus \partial\Omega_h^1$ and $\partial\Omega_h^1 = \{(0, 0), (1, 0), (0, 1), (1, 1)\}$.

For a function U defined on $\bar{\Omega}_h$, the following notation will be used: $U_{ij} = U(x_i, y_j)$, $x_i = ih$, $y_j = jh$, $i, j = 1, 2, \dots, N$ and

$$\begin{aligned} \Delta_x^- U_{ij} &= \frac{U_{ij} - U_{i-1,j}}{h}, & \Delta_x^+ U_{ij} &= \Delta_x^- U_{i+1,j}, \\ \Delta_y^- U_{ij} &= \frac{U_{ij} - U_{i,j-1}}{h}, & \Delta_y^+ U_{ij} &= \Delta_y^- U_{i,j+1}. \end{aligned}$$

1991 AMS Subject Classification 65N12, 65N15

In the linear space of functions defined on $\bar{\Omega}_h$ let

$$[U, V] = h^2 \sum_{i,j=1}^{N-1} U_{ij} V_{ij} + \frac{h^2}{2} \sum_{i=1}^{N-1} (U_{i0} V_{i0} + U_{0i} V_{0i} + U_{Ni} V_{Ni} + U_{iN} V_{iN}) + \frac{h^2}{4} (U_{00} V_{00} + U_{N0} V_{N0} + U_{0N} V_{0N} + U_{NN} V_{NN})$$

be the scalar product and $\|U\| = \sqrt{[U, U]}$ the corresponding norm. The discrete H^1 norm $\|\cdot\|_{1,h}$ is defined by $\|U\|_{1,h} = \sqrt{\|U\|^2 + |U|_{1,h}^2}$, where $\|\cdot\|_{1,h}$ is the discrete H^1 seminorm:

$$|U|_{1,h} = \sqrt{\|\Delta_x^- U\|_x^2 + \|\Delta_y^- U\|_y^2}, \quad \|U\|_x = \sqrt{(U, U)_x}, \quad \|U\|_y = \sqrt{(U, U)_y},$$

$$(U, V)_x = h^2 \sum_{i=1}^N \sum_{j=0}^N U_{ij} V_{ij} \quad \text{and} \quad (U, V)_y = h^2 \sum_{i=0}^N \sum_{j=1}^N U_{ij} V_{ij}.$$

Let T_i , \bar{T}_i and $\overline{\bar{T}}_i$ ($i = 1, 2$) denote the mollifiers defined by

$$T_1 f(x, y) = \int_{-\frac{1}{2}}^{\frac{1}{2}} f(x + th, y) dt, \quad T_2 f(x, y) = \int_{-\frac{1}{2}}^{\frac{1}{2}} f(x, y + th) dt,$$

$$\bar{T}_1 f(0, y) = 2 \int_0^{\frac{1}{2}} f(th, y) dt, \quad \bar{T}_2 f(x, 0) = 2 \int_0^{\frac{1}{2}} f(x, th) dt,$$

$$\overline{\bar{T}}_1 f(1, y) = 2 \int_{-\frac{1}{2}}^0 f(1 + th, y) dt, \quad \overline{\bar{T}}_2 f(x, 1) = 2 \int_{-\frac{1}{2}}^0 f(x, 1 + th) dt.$$

We approximate problem (1) by the finite-difference scheme

$$\Delta_h U = F \quad \text{in } \bar{\Omega}_h, \quad (2)$$

where $\Delta_h U = \Delta_{h,x} U + \Delta_{h,y} U$,

$$\Delta_{h,x} U = \begin{cases} \frac{2}{h}(\Delta_x^+ U - \sigma U), & i = 0, \quad j = 0, 1, 2, \dots, N \\ \Delta_x^+ \Delta_x^- U, & i = 1, 2, \dots, N-1, \quad j = 0, 1, 2, \dots, N, \\ -\frac{2}{h}(\Delta_x^- U + \sigma U), & i = N, \quad j = 0, 1, 2, \dots, N \end{cases}$$

$F_{ij} = T_1 T_2 f_{ij}$, $F_{0j} = \bar{T}_1 T_2 f_{0j}$, ($i, j = 1, 2, \dots, N-1$), $F_{00} = \overline{\bar{T}}_1 \overline{\bar{T}}_2 f_{00}$, and $\Delta_{h,y} U$, F_{Nj} , F_{i0} , F_{iN} , F_{N0} , F_{0N} , F_{NN} defined analogously.

2. Stability of the scheme

To begin, let us prove two lemmas.

LEMMA 1. *Let U, V denote mesh-functions on $\bar{\Omega}_h$. Then $[\Delta_h U, V] = [U, \Delta_h V]$.*

Proof. Using summation by parts it is easy to prove that

$$\begin{aligned} [\Delta_{h,x}U, V] &= -h^2 \sum_{j=1}^{N-1} \sum_{i=1}^N \Delta_x^- U_{ij} \Delta_x^- V_{ij} - \\ &- \frac{h^2}{2} \sum_{i=1}^N (\Delta_x^- U_{i0} \Delta_x^- V_{i0} + \Delta_x^- U_{iN} \Delta_x^- V_{iN}) - h \sum_{i=1}^{N-1} (\sigma_{Ni} U_{Ni} V_{Ni} + \sigma_{0i} U_{0i} V_{0i}) \\ &- \frac{h}{2} (\sigma_{00} U_{00} V_{00} + \sigma_{0N} U_{0N} V_{0N} + \sigma_{N0} U_{N0} V_{N0} + \sigma_{NN} U_{NN} V_{NN}) = [U, \Delta_{h,x}V]. \end{aligned}$$

The operator $\Delta_{h,y}$ has the same property. Therefore,

$$[\Delta_h U, V] = [\Delta_{h,x}U, V] + [\Delta_{h,y}U, V] = [U, \Delta_{h,x}V] + [U, \Delta_{h,y}V] = [U, \Delta_h V]. \quad \blacksquare$$

LEMMA 2. Let U denote mesh-function on $\bar{\Omega}_h$. Then $[\Delta_h U, U] \leq -C|[U]|^2$, where $C = \min\{1, 2\sigma_0\}$.

Proof. For fixed $j = 0, 1, 2, \dots, N$, using an inequality from [8] we get

$$\max\{U_{ij}^2 : 0 \leq i \leq N\} \leq 2h \sum_{i=1}^N (\Delta_x^- U_{ij})^2 + U_{0j}^2 + U_{Nj}^2.$$

This yields the following inequality:

$$h \sum_{i=1}^{N-1} U_{ij}^2 + \frac{h}{2} (U_{0j}^2 + U_{Nj}^2) \leq 2h \sum_{i=1}^N (\Delta_x^- U_{ij})^2 + U_{0j}^2 + U_{Nj}^2.$$

Now let us prove that $[\Delta_{h,x}U, U] \leq -\frac{C}{2}|[U]|^2$. Summing by parts and using last inequality, we obtain:

$$\begin{aligned} [\Delta_{h,x}U, U] &= -h^2 \sum_{j=1}^{N-1} \sum_{i=1}^N (\Delta_x^- U_{ij})^2 - \frac{h^2}{2} \sum_{i=1}^N [(\Delta_x^- U_{i0})^2 + (\Delta_x^- U_{iN})^2] - \\ &- h \sum_{j=1}^{N-1} (\sigma_{Nj} U_{Nj}^2 + \sigma_{0j} U_{0j}^2) - \frac{h}{2} (\sigma_{00} U_{00}^2 + \sigma_{0N} U_{0N}^2 + \sigma_{N0} U_{N0}^2 + \sigma_{NN} U_{NN}^2) \leq \\ &- h \sum_{j=1}^{N-1} \left[h \sum_{i=1}^N (\Delta_x^- U_{ij})^2 + \sigma_0 U_{Nj}^2 + \sigma_0 U_{0j}^2 \right] - \frac{h}{2} \left[h \sum_{i=1}^N (\Delta_x^- U_{i0})^2 + \sigma_0 U_{00}^2 + \sigma_0 U_{N0}^2 \right] \\ &- \frac{h}{2} \left[h \sum_{i=1}^N (\Delta_x^- U_{iN})^2 + \sigma_0 U_{0N}^2 + \sigma_0 U_{NN}^2 \right] \leq -\frac{C}{2}|[U]|^2. \end{aligned}$$

The inequality $[\Delta_{h,y}U, U] \leq -\frac{C}{2}|[U]|^2$ can be proved analogously and we easily obtain Lemma 2. \blacksquare

THEOREM 1. For any $f \in H^s(\Omega)$, $s \geq 0$, finite-difference scheme (2) has unique solution U . Moreover,

$$\|U\|_{1,h} \leq \sqrt{2 + \frac{1}{C}} \|F\|, \quad (3)$$

where $C = \min \{1, 2\sigma_0\}$.

Proof. The existence and uniqueness of solutions follow from the fact that Δ_h is a self-adjoint and negative definite operator (Lemmas 1. and 2.). Further, (using Lemma 2.) we can prove stability in the norm $\|[\cdot]\|$: $\|U\|^2 \leq \frac{1}{C} [-\Delta_h U, U] = \frac{1}{C} [-F, U] \leq \frac{1}{C} \|F\| \|U\|$ and thus $\|U\| \leq \frac{1}{C} \|F\|$. Summing by parts we can also prove that $[\Delta_{h,x} U, U] \leq -\frac{1}{2} \|\Delta_x^- U\|_x^2$, $[\Delta_{h,y} U, U] \leq -\frac{1}{2} \|\Delta_y^- U\|_y^2$, and $\|U\|_{1,h}^2 \leq 2[-\Delta_h U, U]$. Thence and using Lemma 2. we get (3). ■

3. The estimates of energy norm $[-\Delta_h U, U]$

In this section we present three lemmas. Each of them will be used to obtain an appropriate error estimate for scheme (2).

LEMMA 3. Let U denote a mesh-function on $\bar{\Omega}_h$ which is a solution of finite-difference scheme (2). Then

$$[-\Delta_h U, U] \leq C_3 \left\{ h^2 \sum_{i,j=1}^{N-1} F_{ij}^2 + \frac{h^3}{4} \left[\sum_{i=0}^N (F_{i0}^2 + F_{iN}^2) + \sum_{i=1}^{N-1} (F_{0i}^2 + F_{Ni}^2) \right] \right\}, \quad (4)$$

where C_3 is a positive constant.

Proof. Using the ε -inequality: $|ab| \leq \varepsilon a^2 + \frac{1}{4\varepsilon} b^2$; $a, b \in \mathbf{R}$, $\varepsilon > 0$; in the identity $[-\Delta_h U, U] = [-F, U]$ as follows:

$$\begin{aligned} -h^2 \sum_{i,j=1}^{N-1} F_{ij} U_{ij} &\leq \varepsilon h^2 \sum_{i,j=1}^{N-1} U_{ij}^2 + \frac{h^2}{4\varepsilon} \sum_{i,j=1}^{N-1} F_{ij}^2, \\ -\frac{h^2}{2} \sum_{i=1}^{N-1} F_{i0} U_{i0} &\leq \varepsilon h \sum_{i=1}^{N-1} U_{i0}^2 + \frac{h^3}{16\varepsilon} \sum_{i=1}^{N-1} F_{i0}^2 \quad \text{and} \quad -\frac{h^2}{4} F_{00} U_{00} \leq \frac{\varepsilon h}{4} U_{00}^2 + \frac{h^3}{16\varepsilon} F_{00}^2 \end{aligned}$$

we obtain the following inequality:

$$[-\Delta_h U, U] \leq \varepsilon \mathbb{U} + \frac{1}{4\varepsilon} \left\{ h^2 \sum_{i,j=1}^{N-1} F_{ij}^2 + \frac{h^3}{4} \sum_{i=0}^N (F_{i0}^2 + F_{iN}^2) + \frac{h^3}{4} \sum_{i=1}^{N-1} (F_{0i}^2 + F_{Ni}^2) \right\}$$

where $\mathbb{U} = \mathbb{U}(U, h)$, more precisely

$$\mathbb{U} = h^2 \sum_{i,j=1}^{N-1} U_{ij}^2 + h \sum_{i=1}^{N-1} (U_{0i}^2 + U_{i0}^2 + U_{Ni}^2 + U_{iN}^2) + \frac{h}{4} (U_{00}^2 + U_{0N}^2 + U_{N0}^2 + U_{NN}^2).$$

It is easy to prove that $\mathbb{U}(U, h) \leq \left(\frac{1}{C} + \frac{1}{\sigma_0}\right) [-\Delta_h U, U]$, where $C = \min\{1, 2\sigma_0\}$ and $\sigma \geq \sigma_0 > 0$. Thus we get (4) where $C_3 = [4\varepsilon - 4\varepsilon^2 (C^{-1} + \sigma_0^{-1})]^{-1}$ and the value of ε can be chosen so that $1 > \varepsilon (C^{-1} + \sigma_0^{-1})$, the optimal choice being

$$\varepsilon = C\sigma_0(2\sigma_0 + 2C)^{-1} = \begin{cases} 1 + \sigma_0^{-1}, & \sigma_0 \geq \frac{1}{2}, \\ \frac{3}{2}\sigma_0^{-1}, & 0 < \sigma_0 < \frac{1}{2}. \end{cases} \quad \blacksquare$$

Using the same technique, we can prove the following two lemmas. Their proofs are omitted.

LEMMA 4. *Let U denote a mesh-function on $\overline{\Omega}_h$ which is the solution of finite-difference scheme (2). If we substitute F_{ij} by $F_{ij} = \Delta_x^+ \xi_{1,ij} + \Delta_y^+ \xi_{2,ij}$, ($i, j = 1, 2, \dots, N-1$), $F_{i0} = \Delta_x^+ \xi_{1,i0} + \eta_{2,i0}$, $F_{0i} = \eta_{1,0i} + \Delta_y^+ \xi_{2,0i}$, $F_{Ni} = \eta_{1,Ni} + \Delta_y^+ \xi_{2,Ni}$, $F_{iN} = \Delta_x^+ \xi_{1,iN} + \eta_{2,iN}$, ($i = 1, 2, \dots, N-1$), $F_{00} = \eta_{1,00} + \eta_{2,00}$, $F_{0N} = \eta_{1,0N} + \eta_{2,0N}$, $F_{N0} = \eta_{1,N0} + \eta_{2,N0}$ and $F_{NN} = \eta_{1,NN} + \eta_{2,NN}$ then*

$$[-\Delta_h U, U] \leq C_4 \left\{ h^2 \sum_{j=0}^N \sum_{i=1}^N (\xi_{1,ij}^2 + \xi_{2,ji}^2) + h \sum_{i=1}^N (\xi_{1,Ni}^2 + \xi_{1,1i}^2 + \xi_{2,iN}^2 + \xi_{2,i1}^2) + h^3 \sum_{i=0}^N (\eta_{1,i0}^2 + \eta_{1,iN}^2 + \eta_{2,0i}^2 + \eta_{2,Ni}^2) \right\}.$$

LEMMA 5. *Under the same assumptions as in Lemma 4, and defining $\alpha_{0i} = \xi_{1,1i} - \frac{h}{2}\eta_{1,0i}$, $\alpha_{Ni} = -\xi_{1,Ni} - \frac{h}{2}\eta_{1,Ni}$, $\beta_{i0} = \xi_{2,i1} - \frac{h}{2}\eta_{2,i0}$ and $\beta_{iN} = -\xi_{2,iN} - \frac{h}{2}\eta_{2,iN}$, ($i = 0, 1, 2, \dots, N$), the following inequality holds:*

$$[-\Delta_h U, U] \leq C_5 \left\{ h^2 \sum_{j=0}^N \sum_{i=1}^N (\xi_{1,ij}^2 + \xi_{2,ji}^2) + h \sum_{i=0}^N (\alpha_{0i}^2 + \alpha_{Ni}^2 + \beta_{i0}^2 + \beta_{iN}^2) \right\}.$$

4. Convergence of the finite-difference scheme

Before stating our main results we quote the following theorem which is a variant of the well-known Dupont-Scott approximation theorem (see [2]).

THEOREM 2. *Let E be a bounded connected domain in \mathbf{R}^2 satisfying the cone condition and $\mathcal{A}(u)$ a bounded linear functional on $H^s(E)$ ($s = \{s\} + \alpha$, $\{s\} \geq 0$ is integer and $\{s\} < s$, $0 < \alpha \leq 1$) such that $P_{\{s\}} \subset \text{Kernel}(\mathcal{A}(u))$, where $P_{\{s\}}$ denotes the set of polynomials of degree $\leq \{s\}$. Then, for any $u \in H^s(E)$, $|\mathcal{A}(u)| \leq C|u|_{H^s(E)}$, where $C = C(E, s)$ is a positive constant independent of u and $|\cdot|_{H^s(E)}$ is the highest seminorm of $H^s(E)$.*

The derivations of all error estimates below are based on the above theorem.

THEOREM 3. *Suppose that $u \in H^s(\Omega)$, $2 \leq s \leq 4$, is the solution of problem (1) and U is the solution of the finite-difference scheme (2). Then*

$$\| [U - u] \|_{1,h} \leq Ch^{s-2} \| u \|_{H^s(\Omega)} = O(h^{s-2}).$$

Proof. Let us define the global error as $z = U - u$. Then $\Delta_h z_{ij} = \Delta_h U_{ij} - \Delta_h u_{ij} = F_{ij} - \Delta_h u_{ij} = \varphi_{ij}$. We shall consider three cases:

i) If $(ih, jh) \in \Omega_h$, then

$$\varphi_{ij} = T_2 \Delta_x^- \frac{\partial u}{\partial x} \left(ih + \frac{h}{2}, jh \right) + T_1 \Delta_y^- \frac{\partial u}{\partial x} \left(ih, jh + \frac{h}{2} \right) - \Delta_h u_{ij}.$$

Using Theorem 2 and standard technique based on Theorem 2 as in [3], [5] or [9], we obtain $|\varphi_{ij}| \leq Ch^{s-3} |u|_{H^s(e_{ij})}$, $2 < s \leq 4$, where $e_{ij} = \{(x, y) : ih - h \leq x \leq ih + h; jh - h \leq y \leq jh + h\}$.

ii) If $(ih, jh) \in \partial\Omega_h^2$, for example $(0, jh)$, then

$$\begin{aligned} \varphi_{0j} &= \frac{2}{h} T_2 \left[\frac{\partial u}{\partial x} \left(\frac{h}{2}, jh \right) - \frac{\partial u}{\partial x} (0, jh) \right] + \bar{T}_1 \Delta_y^- \frac{\partial u}{\partial y} \left(0, jh + \frac{h}{2} \right) - \\ &\quad - \frac{1}{h^2} \left[u_{0,j+1} + u_{0,j-1} + 2u_{1,j} - 2h \frac{\partial u}{\partial x} (0, jh) - 4u_{0j} \right]. \end{aligned}$$

In the same way, except that $2 < s \leq 3$, we obtain $|\varphi_{0j}| \leq Ch^{s-3} |u|_{H^s(e_{0j})}$, where $e_{0j} = \{(x, y) : 0 \leq x \leq h; jh - h \leq y \leq jh + h\}$.

iii) If $(ih, jh) \in \partial\Omega_h^1$, for example $(0, 0)$, then

$$\begin{aligned} \varphi_{00} &= \frac{2}{h} \bar{T}_2 \left[\frac{\partial u}{\partial x} \left(\frac{h}{2}, 0 \right) - \frac{\partial u}{\partial x} (0, 0) \right] + \frac{2}{h} \bar{T}_1 \left[\frac{\partial u}{\partial y} \left(0, \frac{h}{2} \right) - \frac{\partial u}{\partial y} (0, 0) \right] - \\ &\quad - \frac{2}{h^2} \left[u_{10} - u_{00} - h \frac{\partial u}{\partial x} (0, 0) + u_{01} - u_{00} - h \frac{\partial u}{\partial y} (0, 0) \right] \end{aligned}$$

and we obtain, provided $2 < s \leq 3$, $|\varphi_{00}| \leq Ch^{s-3} |u|_{H^s(e_{00})}$, where $e_{00} = \{(x, y) : 0 \leq x, y \leq h\}$.

However, we can obtain $\| [U] \|_{1,h}^2 \leq C [\Delta_h U, U]$. Thence, using (4), we get

$$\| [z] \|_{1,h}^2 \leq C \left\{ h^2 \sum_{i,j=1}^{N-1} \varphi_{ij}^2 + \frac{h^3}{4} \sum_{i=0}^N (\varphi_{i0}^2 + \varphi_{iN}^2) + \frac{h^3}{4} \sum_{i=1}^{N-1} (\varphi_{0i}^2 + \varphi_{Ni}^2) \right\}.$$

Now, for $2 < s \leq 3$, it is easy to prove that $\| [z] \|_{1,h} \leq Ch^{s-2} |u|_{H^s(\Omega)}$. If $3 < s \leq 4$, then

$$h^2 \sum_{i,j=1}^{N-1} \varphi_{ij}^2 \leq Ch^{2s-4} |u|_{H^s(\Omega)}^2. \quad (5)$$

On the other hand, if $u \in H^s(\Omega)$, then $u \in H^3(\Omega)$ and

$$\frac{h^3}{4} \left[\sum_{i=0}^N (\varphi_{i0}^2 + \varphi_{iN}^2) + \sum_{i=1}^{N-1} (\varphi_{0i}^2 + \varphi_{Ni}^2) \right] \leq Ch^3 |u|_{H^3(\partial_h \Omega)}, \quad (6)$$

where $\partial_h \Omega$ is the boundary strip of width h . But, according to [7]

$$|u|_{H^3(\partial_h \Omega)} \leq C \|u\|_{H^s(\Omega)} \cdot \begin{cases} h^{s-3}, & 3 < s < \frac{7}{2} \\ \sqrt{h} |\ln h|, & s = \frac{7}{2} \\ \sqrt{h}, & \frac{7}{2} < s \leq 4. \end{cases} \quad (7)$$

Using (5), (6) and (7) we obtain $\|z\|_{1,h} \leq Ch^{s-2} \|u\|_{H^s(\Omega)}$ and that completes the proof. ■

THEOREM 4. *Suppose that $u \in H^s(\Omega)$, $\frac{3}{2} < s \leq 3$, is the solution of problem (1) where $\sigma \in M(H^{s-1}(0,1))$ (see [6]) and U is the solution of (2). Then*

$$\|U - u\|_{1,h} = \begin{cases} O(h^{s-1}), & \frac{3}{2} < s < \frac{5}{2} \\ O(h\sqrt{h} |\ln h|), & s = \frac{5}{2} \\ O(h\sqrt{h}), & \frac{5}{2} < s \leq 3. \end{cases}$$

Proof. This theorem is similar to the previous one. Therefore we begin the proof as before. Naturally, this time we shall use Lemma 5 and thus we have to derive the following:

i) If $i = 1, 2, \dots, N$; $j = 1, 2, \dots, N-1$ and $\frac{3}{2} < s \leq 3$, then

$$\xi_{1,ij} = T_2 \left[\frac{\partial u}{\partial x} \left(ih - \frac{h}{2}, jh \right) \right] - \Delta_x^- u_{ij} \quad \text{and} \quad |\xi_{1,ij}| \leq Ch^{s-2} |u|_{H^s(e_{ij})},$$

where $e_{ij} = \{(x, y) : ih - h \leq x \leq ih, jh - \frac{h}{2} \leq y \leq jh + \frac{h}{2}\}$.

ii) If $i = 1, 2, \dots, N$; $j = 0$ or $j = N$ and $\frac{3}{2} < s \leq 2$, then

$$\xi_{1,i0} = \overline{T}_2 \left[\frac{\partial u}{\partial x} \left(ih - \frac{h}{2}, 0 \right) \right] - \Delta_x^- u_{i0}, \quad \xi_{1,iN} = \overline{\overline{T}}_2 \left[\frac{\partial u}{\partial x} \left(ih - \frac{h}{2}, 1 \right) \right] - \Delta_x^- u_{iN}$$

$$\text{and} \quad |\xi_{1,ij}| \leq Ch^{s-2} |u|_{H^s(e_{ij})}$$

where $e_{i0} = \{(x, y) : ih - h \leq x \leq ih, 0 \leq y \leq \frac{h}{2}\}$

or $e_{iN} = \{(x, y) : ih - h \leq x \leq ih, 1 - \frac{h}{2} \leq y \leq 1\}$.

(Analogous results can be obtained for ξ_2 .)

iii) If $j = 1, 2, \dots, N-1$, then $\alpha_{0j} = T_2(\sigma_{0j} u_{0j}) - \sigma_{0j} u_{0j}$ and

$$|\alpha_{0j}| \leq Ch^{s-\frac{3}{2}} |\sigma u|_{H^{s-1}(d_{0j})}, \quad 1 \leq j \leq N,$$

where $d_{0j} = [jh - \frac{h}{2}, jh + \frac{h}{2}]$. (The same results can be obtained for α_{Nj} , β_{jN} and β_{j0} .)

iv) If $i = 0$ and $j = 0$, then $\alpha_{00} = \overline{T}_2(\sigma_{00}u_{00}) - \sigma_{00}u_{00}$ and

$$|\alpha_{00}| \leq Ch^{s-\frac{3}{2}}|\sigma u|_{H^{s-1}(d_{00})}, \quad 1 \leq s \leq 2,$$

where $d_{00} = [0, \frac{h}{2}]$. (The same results can be obtained for α_{ij} and β_{ij} where $(ih, jh) \in \partial\Omega_{h^1}$.)

Thence:

$$h^2 \sum_{i=1}^N \sum_{j=1}^{N-1} (\xi_{1,ij}^2 + \xi_{2,ji}^2) \leq Ch^{2s-2}|u|_{H^s(\Omega)}^2, \quad \frac{3}{2} < s \leq 3,$$

$$h^2 \sum_{i=1}^N (\xi_{1,i0}^2 + \xi_{1,iN}^2 + \xi_{2,0i}^2 + \xi_{2,Ni}^2) \leq C\|u\|_{H^s(\Omega)}^2 \cdot \begin{cases} h^s, & \frac{3}{2} < s < \frac{5}{2} \\ h^3 \ln^2 h, & s = \frac{5}{2} \\ h^3, & \frac{5}{2} < s \leq 3 \end{cases},$$

$$h \sum_{j=1}^{N-1} \alpha_{0j}^2 \leq Ch^{2s-2}|\sigma u|_{H^{s-1}(0,1)}^2, \quad 1 \leq s \leq 3,$$

$$h\alpha_{00}^2 \leq C\|\sigma u\|_{H^{s-1}(0,1)}^2 \cdot \begin{cases} h^{2s-2}, & 1 \leq s \leq \frac{5}{2} \\ h^3 \ln^2 h, & s = \frac{5}{2} \\ h^3, & \frac{5}{2} < s \leq 3 \end{cases}$$

and

$$\begin{aligned} |\sigma u|_{H^{s-1}(0,1)} &\leq \|\sigma u\|_{H^{s-1}(0,1)} \leq C\|\sigma\|_{M(H^{s-1}(0,1))}\|u\|_{H^{s-1}(0,1)} \leq \\ &\leq C\|\sigma\|_{M(H^{s-1}(0,1))}\|u\|_{H^s(\Omega)}. \end{aligned}$$

Now using Lemma 5, we easily complete the proof of the theorem. ■

REFERENCES

- [1] R. A. Adams, *Sobolev Spaces*, Academic Press, New York 1975
- [2] T. Dupont and R. Scott, *Polynomial approximation of functions in Sobolev spaces*, Math. Comp. **34** (1980), 441–463
- [3] B. S. Jovanović, *On the convergence of discrete solutions to the generalized solutions of boundary value problems*, in: *Variational-Difference Methods in Mathematical Physics*, (N. S. Bahvalov, ed.), Academy of Sciences of the USSR, Moscow 1984, 120–129 (in Russian)
- [4] O. A. Ladizhenskaya, *Boundary Value Problems of Mathematical Physics*, Nauka, Moscow 1973 (in Russian)
- [5] R. D. Lazarov, *On the question of convergence of finite difference schemes for generalized solutions of the Poisson equation*, Diff. Uravneniya **17** (1981), 1287–1294 (in Russian)
- [6] V. G. Maz'ya and T. O. Shaposhnikova, *Theory of Multipliers in Spaces of Differentiable Functions*, Monographs Stud. Math. **23**, Pitman, Boston 1985
- [7] L. A. Oganessian and L. A. Ruhovec, *Variational-Difference Methods of Solving Elliptic Equations*, A.N. ArmSSR, Erevan 1979 (in Russian)
- [8] A. A. Samarskiĭ, *Theory of Difference Schemes*, Nauka, Moscow 1983 (in Russian)
- [9] E. E. Süli, B. S. Jovanović and L. D. Ivanović, *Finite difference approximations of generalized solutions*, Math. Comp. **45** (1985), 319–327

(received 08.05.1995.)

University of Kragujevac, Faculty of Sciences

Radoja Domanovića 12, 34000 Kragujevac, Yugoslavia